

Nijenhuis tensors in pseudoholomorphic curves neighborhoods

Boris S. Kruglikov

Abstract

We define and study normal bundle to a pseudoholomorphic curve. Normal forms of 1-jets of the almost complex structure along the curve are deduced via the Nijenhuis tensors distribution. We study the problem of pseudoholomorphic foliation of the curve neighborhood and get some results on persistence of pseudoholomorphic tori.

Introduction

Let $J : TM \rightarrow TM$ be an almost complex structure $J^2 = -\mathbf{1}$. A submanifold $N \subset M$ is called *pseudoholomorphic* (PH-submanifold) if $TN \subset TM$ is J -invariant. Generically the only PH-submanifolds have $\dim_{\mathbb{C}} N = 1$ ([?]).

Local existence of PH-curves was established by Nijenhuis and Woolf ([NW]). The global existence result is due to Gromov ([G]). Non-exceptional PH-spheres usually occur in families ([MS]) and the same concerns PH-disks ([K3]). However even in the complex situation holomorphic tori are generically discrete ([A1]).

In [Mo] Moser posed a question what type of KAM-theory can be constructed for a PH-foliation of an almost complex torus T^{2n} . T^2 -foliations are exceptional: Generically PH-tori are discrete and their number was investigated by Kuksin ([Ku1, Ku2]). Moser considered a foliation by entire PH-lines $\mathbb{C} \rightarrow T^{2n}$ with the slope of general position. The main result states that under a small almost complex perturbations of the standard complex structure J_0 on $T^{2n} = \mathbb{C}^n/\Gamma$, Γ being a cocompact lattice, many leaves persist. If the perturbation is big but tame-restricted then only some of the leaves persist. This was proved by Bangert in [B]. Another proof is given in [KO].

In [A2](1993-25) Arnold asks about almost complex version (in the spirit of the above Moser's result) for his Floquet-type theory of elliptic curves neighborhoods ([A1]). It will be shown the direct extension fails (there are moduli in normal forms), though we conjecture the right generalization is a possibility of PH-foliation of a PH-torus neighborhood by cylinders. We get here partial results and think the general case can be treated by the Moser's method.

The structure of the paper is as follows. In section 1 we review the structure of the Nijenhuis tensor and discuss invariants of almost complex structures and

PH-curves. We recall the relevant holomorphic facts as well.

In section 2 we define the canonical almost complex structure on the normal bundle $N_{\mathcal{C}}M = TM/T\mathcal{C}$ to a PH-curve \mathcal{C} and study its properties. Using it we get a normal form of 1-jet of an almost complex structure along a PH-curve.

In section 3 we study foliations, isolated PH-curves and hyperbolicity and recover some traces of normal bundles to holomorphic curves theory in almost complex situation. We also obtain geometric interpretation of the Moser's non-deformation examples from [Mo].

1. Moduli of PH-curves neighborhoods

Throughout this section we assume $\dim M = 4$.

1.1. Nijenhuis tensor characteristic distribution

Nijenhuis tensor of an almost complex structure J is the $(2, 1)$ -tensor

$$N_J \in \Lambda^2 T^*M \otimes TM, \quad N_J(\xi, \eta) = [J\xi, J\eta] - J[J\xi, \eta] - J[\xi, J\eta] - [\xi, \eta]. \quad (1)$$

Integrability of J is expressed via it as $N_J = 0$ ([NW]).

This tensor satisfies the property $N_J(J\xi, \eta) = N_J(\xi, J\eta) = -JN_J(\xi, \eta)$ and so can be considered as an antilinear map $N_J : \Lambda^2 \mathbb{C}^2 \rightarrow \mathbb{C}^2$, $\mathbb{C}^2 = (T_x M^4, J)$. Thus the image is invariant under J and if $N_J \neq 0$ it is a complex line $\mathbb{C} \subset \mathbb{C}^2$.

Definition 1. *Let us call $\Pi^2 = \text{Im } N_J \subset TM$ the Nijenhuis tensor characteristic distribution on 4-dimensional almost complex manifold (M^4, J) .*

In fact Π is a differential systems. At singular points, which are generically isolated, the rank of Π is 0.

This distribution Π^2 is in general situation nonintegrable. Therefore it has nontrivial derivative $\Pi^3 = \partial\Pi^2$, that is the differential system with the $C^\infty(M)$ -module of sections $\mathcal{P}_3 = C^\infty(\Pi^3)$ generated by the module of sections $\mathcal{P}_2 = C^\infty(\Pi^2)$ and its self-commutator: $\mathcal{P}_3 = [\mathcal{P}_2, \mathcal{P}_2]$. Π^3 is not distribution everywhere and its singularities form a stratified submanifold Σ'_2 of codim = 2.

Consider a point outside Σ'_2 . If the distribution Π^3 is nonintegrable, then $\mathcal{P}_4 = [\mathcal{P}_3, \mathcal{P}_3] = \mathcal{D}(M)$ is the module of all vector fields, so that the next distribution is the whole tangent space. Moreover generically $[\mathcal{P}_2, \mathcal{P}_3] = \mathcal{D}(M)$ outside another stratified submanifold Σ''_2 of codim = 2.

If $x \notin \Sigma'_2$, then $\Pi_x^2 \subset \Pi_x^3$ has a transversal measure. In fact since the J -antilinear isomorphism $N_J(\cdot, \xi_3) : \Pi_x^2 \rightarrow \Pi_x^2$ is orientation reversing, there exist vectors $\xi_1, \xi_2 \in \Pi_x^2$, $\xi_3 \in \Pi_x^3 \setminus \Pi_x^2$ such that $N_J(\xi_1, \xi_3) = \xi_1$, $N_J(\xi_2, \xi_3) = -\xi_2$. These ξ_1, ξ_2 are defined up to multiplication by a constant, while $\xi_3 \pmod{\Pi_x^2}$ is defined up to multiplication by ± 1 . Therefore Π^3/Π^2 is normed. By a similar reason $T_x M/\Pi_x^3$ is normed outside Σ'_2 via the vector $\xi_4 = J\xi_3$.

Note that Π_x^3/Π_x^2 is oriented. Actually $[\xi_1, \xi_2] \pmod{\Pi_x^2}$ depends only on the values of ξ_1, ξ_2 at the point x . It is a vector $f\xi_3 \pmod{\Pi_x^2}$ for some f . So if

we require $\xi_2 = J\xi_1$ then ξ_3 can be chosen so that $f > 0$. This produces coorientation of $\Pi_x^2 \subset \Pi_x^3$ and then via J coorientation of $\Pi_x^3 \subset T_x M$.

Moreover the requirement $f = 1$ determines canonically vector field ξ_1 (still however up to ± 1) and hence $\xi_2 = J\xi_1$. Then we set $\xi_3 = [\xi_1, \xi_2]$ and $\xi_4 = J\xi_3$. So the pair (ξ_1, ξ_2) is defined canonically up to a sign and the pair (ξ_3, ξ_4) is absolutely canonical. The following statement generalizes theorem 7 [K1]:

Theorem 1. *Let almost complex structure J be of general position. Then at generic points $x \in M^4$ a canonical frame $(\xi_1, \xi_2, \xi_3, \xi_4)$ is defined. It restores uniquely the almost complex operator J and the tensor N_J by the tables:*

X	JX
ξ_1	ξ_2
ξ_2	$-\xi_1$
ξ_3	ξ_4
ξ_4	$-\xi_3$

$N_J(\uparrow, \leftarrow)$	ξ_1	ξ_2	ξ_3	ξ_4
ξ_1	0	0	ξ_1	$-\xi_2$
ξ_2	0	0	$-\xi_2$	$-\xi_1$
ξ_3	$-\xi_1$	ξ_2	0	0
ξ_4	ξ_2	ξ_1	0	0

Note that reducing a geometric structure to a frame ($\{e\}$ -structure) solves completely the equivalence problem. The idea is as follows. Consider the moduli of the problem, i.e. functions c_{jk}^i given by the formula $[\xi_j, \xi_k] = \sum c_{jk}^i \xi_i$. Denote by $\mathbb{A} = \{c_{jk}^i\}$ the space of all invariants and by $\Phi : M \rightarrow \mathbb{A}$ the "momentum map" $x \mapsto \{c_{jk}^i(x)\}$. Then two equivalent structures have the same images and the equivalence follows. See [S] for more details.

Remark 1. *Outside $\Sigma'_2 \cup \Sigma''_2$ the distribution Π^2 has a canonical local normal form and is called Engel distribution. Locally it can be realized as Nijenhuis tensors characteristic distributions ([K2]).*

1.2. Invariants of a PH-curve neighborhood

Let \mathcal{C}^2 be a PH-curve. At every point $x \in \mathcal{C}$ we have two J -invariant planes $T_x \mathcal{C}^2$ and Π_x^2 , which generically intersects by zero, except at a finite number of points $\tilde{\Sigma}_0 \subset \mathcal{C}$. The sets $\Sigma'_0 = \Sigma'_2 \cap \mathcal{C}$ and $\Sigma''_0 = \Sigma''_2 \cap \mathcal{C}$ are generically finite as well. The arrangement of all these points

$$\Sigma_0 = \tilde{\Sigma}_0 \cup \Sigma'_0 \cup \Sigma''_0 \subset \mathcal{C}$$

gives a (finite-dimensional) invariant.

For points $x \in \mathcal{C} \setminus \Sigma'_0$ we define field of directions $L^1 = T\mathcal{C} \cap \Pi^3$. The integral curves of this 1-distribution foliate the set $\mathcal{C} \setminus \Sigma'_0$ and in general \mathcal{C} foliates with only nondegenerate singular points. Denote the number of elliptic points by $e(L^1)$ and the number of hyperbolic by $h(L^1)$. Note that (topologically stable) points of $\tilde{\Sigma}_0$ are usually regular points of L^1 . One can prove:

Lemma 2. *Under C^1 -small perturbation of the structure J the foliation L^1 has minimal number of singularities, $\min\{e(L), h(L)\} = 0$, $\max\{e(L), h(L)\} = \frac{1}{2}|\chi(\mathcal{C})|$. For instance if $\mathcal{C} = T^2$ we get foliation without singularities.*

Due to the previous subsection the foliation L^1 is oriented, cooriented and has parallel and transverse measures outside Σ_0 . Thus there exist canonical vector fields v_1 along L^1 and $v_2 = Jv_1$ transverse to it. Consequently the curve \mathcal{C} has a lot of dynamical invariants like winding classes of v_1 and v_2 , the set of critical points, linearizations at them etc. Moreover decomposing

$$[v_1, v_2] = \gamma_1 v_1 + \gamma_2 v_2.$$

we obtain two invariant (under pseudoholomorphic isomorphisms) functions γ_1, γ_2 . These together with the germs of the functions c_{jk}^i from the previous subsection form *moduli* of the \mathcal{C} -neighborhoods germ. They solve the equivalence problem for PH-embeddings $\mathcal{C}^2 \rightarrow M^4$ (of general position).

Example. Let $M = T^2(\varphi, \psi) \times \mathbb{R}^2(x, y)$ be equipped with the structure

$$\begin{aligned} J\partial_x &= \partial_y; & J\partial_\varphi &= \frac{2-\rho y^2}{2}\partial_\psi + \frac{y^2}{2}\partial_\varphi + x\partial_x; \\ J\partial_y &= -\partial_x; & J\partial_\psi &= \frac{4+y^4}{2\rho y^2-4}\partial_\varphi - \frac{y^2}{2}\partial_\psi + \frac{xy^2}{\rho y^2-2}\partial_x + \frac{2x}{\rho y^2-2}\partial_y, \end{aligned}$$

Then $\mathcal{C} = \{x = y = 0\}$ is a PH-torus and the winding number of v_1 is ρ . Similarly one shows the other considered invariants are non-trivial.

As we will see the moduli space of germs of \mathcal{C} -embeddings is much larger than this for the normal bundles, implying impossibility of equivalence [A2](1993-25).

1.3. Holomorphic and pseudoholomorphic structures in a neighborhood of a curve

For a holomorphic curve \mathcal{C} in a complex manifold M^{2n} its normal bundle is defined via the exact sequence:

$$0 \rightarrow T\mathcal{C} \rightarrow TM|_{\mathcal{C}} \rightarrow N_{\mathcal{C}}M \rightarrow 0. \quad (2)$$

Using charts one can check the total space of the bundle is a complex manifold making $N_{\mathcal{C}}M$ into a holomorphic vector bundle over \mathcal{C} .

Another way to introduce the complex structure in $N_{\mathcal{C}}M$ for embedded \mathcal{C} is to use the "box theorem", stating that some neighborhood \mathcal{O} is a holomorphic bundle over \mathcal{C} with fibers biholomorphic to the ball $B^{2n-2}(\varepsilon) \subset \mathbb{C}^{n-1}$. Let A_t be the dilation along the fibers. For the complex structure J on \mathcal{O} we define:

$$J_t = A_t J A_t^{-1} \text{ and } \hat{J} = \lim_{t \rightarrow \infty} J_t. \quad (3)$$

Then \hat{J} is the required (canonical) complex structure on $N_{\mathcal{C}}M$.

Recall that a rational curve is the Riemann sphere $\mathcal{C} = \mathbb{C} \simeq S^2$ and an elliptic curve is a torus $\mathcal{C} = \mathbb{C}/\Gamma \simeq T^2$. This torus is uniquely characterized by its lattice $\Gamma = \mathbb{Z}^2(2\pi, \nu)$, where the first period is fixed to be 2π and for the second $\text{Im}(\nu) > 0$.

All holomorphic topologically trivial 1-dimensional vector bundles over this elliptic curve are described as follows. Consider \mathbb{C}^2 with coordinates (z, φ) and for $\lambda \in \mathbb{C} \setminus \{0\}$, $\nu \in \mathbb{C} \setminus \mathbb{R}$ make the identifications

$$(z, \varphi) \simeq (z, \varphi + 2\pi) \simeq (\lambda z, \varphi + \nu). \quad (4)$$

The bundle $E \rightarrow T^2$ is now given by $(z, \varphi) \mapsto \varphi$. Similarly one classifies S^2 -bundles with fixed two distinguished holomorphic sections "zero" and "infinity".

Arnold in [A1] §27 considers normal bundle $N_{T^2}M^4$ to an elliptic curve and prove that if the pair (λ, ν) is normal nonresonant ($\lambda^n \neq e^{ik\nu}$ plus some diophantine condition for this pair) then a small neighborhood of $T^2 \subset M$ is biholomorphically equivalent to a neighborhood of the zero section in $N_{T^2}M$.

Note that the number λ is defined by 1-jet of the complex structure on the torus T^2 . But in almost complex situation 1-jet is determined by the field of the Nijenhuis tensors N_J along the torus, so this λ is not defined. Moreover with another definition of the normal bundle proposed we'll see the whole Nijenhuis tensor does matter. For $\dim M = 4$ the situation simplifies due to

Proposition 3. *Small neighborhood \mathcal{O} of a PH-curve $\mathcal{C} \subset M^4$ can be foliated by transversal PH-disks D^2 .*

This actually follows from Nijenhuis-Wolf theorem [NW] of existence of small PH-disk in a given direction (which is normal in our case) and smooth dependence on this direction. Using proposition 3 we can define an almost complex structure \hat{J} on $N_{\mathcal{C}}M$ making it into an almost complex bundle (cf. [M2]) via formula (3). However for almost complex structures this approach works only in dimension 4 and being introduced so the properties of this almost complex structure remains hidden. We will handle the general case via differential-geometric tools in the next section.

2. Differential geometry around PH-curves

In this section we consider the general case $\dim M = 2n$.

2.1. Normal bundle of a pseudoholomorphic curve

In almost complex case the bundle $TM|_{\mathcal{C}}$ for a PH-curve $\mathcal{C} \subset M$ is no longer holomorphic. So one needs to consider almost complex bundles. A bundle $\pi : (E, J) \rightarrow (\mathcal{C}, J_0)$ is called almost complex if

$$\pi_*(J\xi) = J_0(\pi_*\xi).$$

Proposition 4. *Consider the normal bundle to a PH-curve $\mathcal{C} \subset M$ given as vector space by the sequence (2). There is a canonical almost complex structure \hat{J} on $N_{\mathcal{C}}M$ such that $(N_{\mathcal{C}}M, \hat{J})$ is an almost complex bundle.*

Proof. Recall that on almost complex manifold (M, J) there is always an almost complex connection $\nabla J = 0$. Actually if ∇' is any linear connection then $\nabla = \frac{1}{2}(\nabla' - J\nabla'J)$ is also a linear connection, which preserves the structure J . Moreover we can assume that ∇ is minimal, i.e. the torsion of ∇ equals to its antilinear by each argument part $T_\nabla = T_\nabla^- = \frac{1}{4}N_J$ ([L]).

Lemma 5. *There exists a minimal almost complex connection such that the curve \mathcal{C} is a totally geodesic submanifold, i.e. parallel transport of any vector $v \in T\mathcal{C}$ along a path in \mathcal{C} belongs again to $T\mathcal{C}$.*

Proof. To see this let us make a gauge transformation $\tilde{\nabla} = \nabla + A$, where $A \in \Omega^1(M, \text{end}_{\mathbb{C}} \tau_M)$ is a 1-form with values in complex endomorphisms of the tangent bundle. If we require that this 1-form is symmetric $A \in S^2 T^*M \otimes_{\mathbb{C}} TM$, then the new connection $\tilde{\nabla}$ is also almost complex and minimal.

Let ξ be a vector field on \mathcal{C} with nondegenerate critical points, $\xi \in \mathcal{D}(\mathcal{C})$. Let $\nabla_\xi \xi = \eta \in \mathcal{D}(M)$. We define $A(\xi, \xi) = -\eta$, $A(J\xi, \xi) = A(\xi, J\xi) = -J\eta$, $A(J\xi, J\xi) = \eta$ and for other vectors somehow preserving symmetry and J -linearity (near the critical points there's a more work, we need to choose ξ so that η has a zero of the second order at these critical points).

Then $\tilde{\nabla}_\xi \xi = 0$, $\tilde{\nabla}_\xi J\xi = 0$. Therefore by minimality $\tilde{\nabla}_{J\xi} \xi = \tilde{\nabla}_\xi J\xi + [J\xi, \xi] + \frac{1}{4}N_J(J\xi, \xi) = [J\xi, \xi] \in \mathcal{D}(\mathcal{C})$ and also $\tilde{\nabla}_{J\xi} J\xi \in \mathcal{D}(\mathcal{C})$. So $\tilde{\nabla}$ preserves $T\mathcal{C}$. \square

Another way to prove this is to introduce trivial connection in $\mathcal{O}(\mathcal{C}) \simeq \mathcal{C} \times D$ and then to check that procedures of making connection almost complex and then minimal do not destroy the property of \mathcal{C} being totally geodesic. Let us denote this new connection again by the symbol ∇ .

We introduce a connection $\hat{\nabla}$ to the bundle $N_{\mathcal{C}}M$ by means of parallel transport as follows. Let $v = [\theta] \in (N_{\mathcal{C}}M)_x$ be the class of $\theta \in T_x M$ and let $\gamma(t) \subset \mathcal{C}$ be a curve, $\gamma(0) = x$. Calculate parallel transport $\theta(t)$ of θ along $\gamma(t)$. We define $v(t) = [\theta(t)]$ to be the parallel transport of v along $\gamma(t)$. Since \mathcal{C} is totally geodesic, the definition is correct ($\hat{\nabla}$ -parallel transport of 0 is 0). Moreover the connection $\hat{\nabla}$ is \mathbb{R} -linear. So as usual in the theory of generalized connections we conclude that $\hat{\nabla}$ is a linear connection.

Let $a \in N_{\mathcal{C}}M$ be a point in the normal bundle with projection $\pi(a) = x$. Let $T_a(N_{\mathcal{C}}M) = H_a \oplus V_a$ be the decomposition by horizontal and vertical subspaces induced by $\hat{\nabla}$. The first space $H_a \xrightarrow{\pi_*} T_x \mathcal{C}$ has a canonical complex structure J_1 induced from J by π_* , while the second $V_a \simeq T_x M / T_x \mathcal{C}$ inherits canonical complex structure J_2 from J . So we introduce the structure on $N_{\mathcal{C}}M$ by the rule $\hat{J} = J_1 \oplus J_2$.

Let us show that the constructed almost complex structure \hat{J} does not depend on the choice of minimal connection ∇ preserving $T\mathcal{C}$. Let us change the connection $\tilde{\nabla} = \nabla + A$, $A \in \Omega^1(M; \text{end}_{\mathbb{C}} \tau_M) \cap S^2 \tau_M^* \otimes \tau_M$. This affects in the change of decomposition $T_a(N_{\mathcal{C}}M) = \tilde{H}_a \oplus V_a$, where $\tilde{H}_a = \text{graph}\{\lambda_a : H_a \rightarrow V_a\}$ and $\lambda_a = A(\cdot)a$. We state that λ_a is a complex linear map. Actually

$$\lambda_a(Jw) = A(Jw)a = A(a)Jw = JA(a)w = JA(w)a = J\lambda_a(w).$$

So the complex structure \hat{J} on $T_a(N_{\mathcal{C}}M)$ is canonically defined. \square

Remark 2. Let $\phi_t : (\mathcal{C}, J_0) \rightarrow (M^4, J)$ be a family of J -holomorphic curves of the same holomorphic type with $\phi_0 = \text{id}$. Then $\frac{d}{dt}\big|_{t=0} \phi_t$ is the pseudoholomorphic curve in $N_{\mathcal{C}}M$. Moreover we have a one-parametric (scaled) family of PH-curves. Since $(N_{\mathcal{C}}M)_x = T_x M / T_x \mathcal{C}$ this family does not depend on the reparametrization $\tilde{\phi}_t = \phi_t \circ g_t$ for $g_t \in \text{Aut}(\mathcal{C}, J_0)$ (note that a curve in the bundle bijectively projected to the base has a natural parametrization). We will use it in §3.4.

2.2 Almost complex bundles

Now we return to the case $\dim M = 4$.

Proposition 6. Let $\pi : (E^4, J) \rightarrow (\mathcal{C}^2, J_0)$ be an almost complex bundle over a curve. Then the characteristic distribution $\Pi^2 = \text{Im } N_J$ is integrable and is tangent to the fibers $F_x = \pi^{-1}(x)$.

Proof. Actually:

$$\begin{aligned} \pi_* N_J(\xi, \eta) &= \pi_*[J\xi, J\eta] - \pi_*J[\xi, J\eta] - \pi_*J[J\xi, \eta] - \pi_*[\xi, \eta] \\ &= [J_0\pi_*\xi, J_0\pi_*\eta] - J_0[\pi_*\xi, J_0\pi_*\eta] - J_0[J_0\pi_*\xi, \pi_*\eta] - [\pi_*\xi, \pi_*\eta] \\ &= N_{J_0}(\pi_*\xi, \pi_*\eta) = 0. \quad \square \end{aligned}$$

Corollary 7. Codimension of the set of almost complex structures, the germs of which on the PH-curve $\mathcal{C} \subset M$ are isomorphic to these of the normal bundle $\mathcal{C} \subset N_{\mathcal{C}}M$, in the set of all almost complex structures is infinity. \square

Proof. Actually if the distribution Π^2 is nonintegrable in a neighborhood of the PH-curve \mathcal{C} , then a neighborhood (\mathcal{O}, J) of the curve is not isomorphic to a neighborhood of the zero section in the normal bundle $(N_{\mathcal{C}}M, \hat{J})$. \square

This property is contrary to its analog in the complex category, see Arnold theorem [A1] about neighborhoods of elliptic curves (§1.3). So extending the category the discussed property becomes exceptional.

As the following example shows the integrability of Π^2 is necessary but by no means sufficient condition on the structure J to be locally isomorphic to its representative on the normal bundle.

Example. Consider the almost complex structure given on a $T^2(\varphi)D^2(z)$ with $\varphi = \varphi_1 + i\varphi_2$, $z = x + iy$ by the formula:

$$J\partial_x = \partial_y, \quad J\partial_{\varphi_1} = \partial_{\varphi_2} + A_1\partial_{\varphi_1} + A_2\partial_{\varphi_2} + B_1\partial_x + B_2\partial_y, \quad (5)$$

with $A_i|_{T^2} = B_i|_{T^2} = 0$. The condition $\Pi^2 = T_*\{\varphi = \text{const}\}$ is equivalent to the following PDE system

$$\begin{cases} \frac{\partial A_1}{\partial y} = A_1 \frac{\partial A_1}{\partial x} - \frac{1 + A_1^2}{1 + A_2} \frac{\partial A_2}{\partial x} \\ \frac{\partial A_2}{\partial y} = (1 + A_2) \frac{\partial A_1}{\partial x} - A_1 \frac{\partial A_2}{\partial x} \end{cases} \quad (6)$$

If the functions A_i satisfy this (Cauchy-Kovalevsky) system the projection along the leaves is given by the formula $(z, \varphi) \mapsto \varphi$. Therefore our structure J is projectable iff $A_i \equiv 0$. But there are nonzero solutions of (6). For example:

$$A_1 = -\frac{x}{1+y}, \quad A_2 = -\frac{y}{1+y}.$$

So the space of projectable structures J are of codim = ∞ among the structures with Π^2 integrable, which are of codim = ∞ among all almost complex structures with a fixed 0-jet on the torus T^2 .

2.3 Nijenhuis tensor of normal bundles

Let $N_{\mathcal{C}}M$ be a normal bundle of pseudoholomorphic curve $\mathcal{C} \subset M^4$. Then at the points $x \in \mathcal{C}$ two different Nijenhuis tensors N_J of ambient structure and $N_{\hat{J}}$ of the structure of the normal bundle are defined. Are there some relations between these two tensors? Or between characteristic distributions of these structures? As the following examples show the answer is negative.

Example. "Parallel distribution Π^2 ". Let $M = \mathbb{R}^4(x_1, y_1, x_2, y_2)$ be equipped with almost complex structure

$$J\partial_{x_1} = \partial_{y_1}, \quad J\partial_{y_1} = -\partial_{x_1}, \quad J\partial_{x_2} = \partial_{y_2} + x_1\partial_{x_1}, \quad J\partial_{y_2} = -\partial_{x_2} - x_1\partial_{y_1}.$$

Then the curve $\mathcal{C} = \{x_2 = y_2 = 0\}$ is pseudoholomorphic. We calculate that the following are the Nijenhuis tensor of the structure J and a minimal almost complex connection (note that $\Pi^2 = \text{Im } N_J = \langle \partial_{x_1}, \partial_{y_1} \rangle$):

$N_J(\uparrow, \leftarrow)$	∂_{x_1}	∂_{y_1}	∂_{x_2}	∂_{y_2}	$\nabla_{\uparrow} \leftarrow$	∂_{x_1}	∂_{y_1}	∂_{x_2}	∂_{y_2}
∂_{x_1}	0	0	$-\partial_{y_1}$	$-\partial_{x_1}$	∂_{x_1}	0	0	$-\frac{1}{4}\partial_{y_1}$	$-\frac{3}{4}\partial_{x_1}$
∂_{y_1}	0	0	$-\partial_{x_1}$	∂_{y_1}	∂_{y_1}	0	0	$-\frac{1}{4}\partial_{x_1}$	$-\frac{1}{4}\partial_{y_1}$
∂_{x_2}	∂_{y_1}	∂_{x_1}	0	$-x_1\partial_{y_1}$	∂_{x_2}	0	0	0	0
∂_{y_2}	∂_{x_1}	$-\partial_{y_1}$	$x_1\partial_{y_1}$	0	∂_{y_2}	$-\frac{1}{2}\partial_{x_1}$	$-\frac{1}{2}\partial_{y_1}$	$\frac{1}{4}x_1\partial_{y_1}$	$\frac{1}{4}x_1\partial_{x_1}$

So we find that the horizontal planes are $H = \langle \partial_{x_1}, \partial_{y_1} \rangle$, whence the structure on $N_{\mathcal{C}}M$ is

$$\hat{J}\partial_{x_1} = \partial_{y_1}, \quad \hat{J}\partial_{y_1} = -\partial_{x_1}, \quad \hat{J}\partial_{x_2} = \partial_{y_2}, \quad \hat{J}\partial_{y_2} = -\partial_{x_2}$$

and $N_{\hat{J}} = 0$.

Example. "Transversal distribution Π^2 ". Let the structure be now

$$J\partial_{x_1} = \partial_{y_1} + x_2\partial_{x_2}, \quad J\partial_{y_1} = -\partial_{x_1} - x_2\partial_{y_2}, \quad J\partial_{x_2} = \partial_{y_2}, \quad J\partial_{y_2} = -\partial_{x_2}.$$

Again the curve $\mathcal{C} = \{x_2 = y_2 = 0\}$ is pseudoholomorphic and the Nijenhuis tensor and a minimal almost complex connection are (now $\Pi^2 = \langle \partial_{x_2}, \partial_{y_2} \rangle$):

$N_J(\uparrow, \leftarrow)$	∂_{x_1}	∂_{y_1}	∂_{x_2}	∂_{y_2}	∇_{\leftarrow}	∂_{x_1}	∂_{y_1}	∂_{x_2}	∂_{y_2}
∂_{x_1}	0	$-x_2\partial_{y_2}$	∂_{y_2}	∂_{x_2}	∂_{x_1}	0	0	0	0
∂_{y_1}	$x_2\partial_{y_2}$	0	∂_{x_2}	$-\partial_{y_2}$	∂_{y_1}	$\frac{1}{4}x_2\partial_{y_2}$	$\frac{1}{4}x_2\partial_{x_2}$	$-\frac{1}{2}\partial_{x_2}$	$-\frac{1}{2}\partial_{y_2}$
∂_{x_2}	$-\partial_{y_2}$	$-\partial_{x_2}$	0	0	∂_{x_2}	$-\frac{1}{4}\partial_{y_2}$	$-\frac{3}{4}\partial_{x_2}$	0	0
∂_{y_2}	$-\partial_{x_2}$	∂_{y_2}	0	0	∂_{y_2}	$-\frac{1}{4}\partial_{x_2}$	$-\frac{1}{4}\partial_{y_2}$	0	0

The horizontal planes are $H = \langle \partial_{x_1}, \partial_{y_1} + \frac{1}{2}e^{\frac{1}{2}y_1}(x_2\partial_{x_2} + y_2\partial_{y_2}) \rangle$. So the structure on $N_{\mathcal{C}}M$ is

$$\begin{aligned}\hat{J}\partial_{x_1} &= \partial_{y_1} + \frac{1}{2}e^{\frac{1}{2}y_1}(x_2\partial_{x_2} + y_2\partial_{y_2}), & \hat{J}\partial_{x_2} &= \partial_{y_2}, \\ \hat{J}\partial_{y_1} &= -\partial_{x_1} - \frac{1}{2}e^{\frac{1}{2}y_1}(x_2\partial_{y_2} - y_2\partial_{x_2}), & \hat{J}\partial_{y_2} &= -\partial_{x_2}.\end{aligned}$$

Now $N_{\hat{J}}(\partial_{x_1}, \partial_{x_2}) = -e^{\frac{1}{2}y_1}\partial_{y_2}$ and so the characteristic distribution of the normal structure is "the same" as for the structure J : $\hat{\Pi}^2 = \text{Im } N_{\hat{J}} = \langle \partial_{x_2}, \partial_{y_2} \rangle$.

So the answer to the above question is positive if N_J is of special type.

2.4 Linear bundle almost complex structures

Consider an almost complex vector bundle $\pi : (E, J) \xrightarrow{F} (\mathcal{C}, J_0)$ of the rank $\dim F = 2n$ and suppose the restriction $J|_F$ is a linear complex structure on the fiber. So we can also consider (E, π, \mathcal{C}) just as vector bundle with complex structures in the fibers.

Definition 2. We call the almost complex structure J on E linear bundle structure if there exists a linear minimal almost complex connection $\hat{\nabla}$ on this bundle such that the lift $T_b E \xleftarrow{\hat{\nabla}} T_a \mathcal{C}$ is a complex mapping, splitting the exact sequence

$$0 \rightarrow F \rightarrow T_a E \rightarrow T_x \mathcal{C} \rightarrow 0, \quad x = \pi(a).$$

In particular the zero section $\mathcal{C} \subset E$ is a J -pseudoholomorphic curve.

We note now that the almost complex structure \hat{J} on the normal bundle $N_{\mathcal{C}}M$ is a linear bundle structure.

Lemma 8. The Nijenhuis tensor of linear bundle almost complex structure J is constant along the fibers and determines J -invariant differential system $\Pi = \text{Im } N_J \subset TE$ which is a subsystem of vertical distribution F .

Proof. There are local coordinates (φ, z) on $\pi^{-1}(U) = UF$, $\varphi = \varphi_1 + i\varphi_2$, $z_k = x_k + iy_k$, $1 \leq k \leq n$, such that $\hat{\nabla}$ -lift of ∂_{φ_i} is $\partial_{\varphi_i} + \sum b_{ij}\partial_{x_j} + c_{ij}\partial_{y_j}$, where the coefficients are linear functions of z_k and the vertical coordinates are complex linear coordinates but horizontal coordinates are complex only on the zero section. So the structure J is given by relations

$$J\partial_{\varphi_1} = \partial_{\varphi_2} + \sum (b_{2j} + c_{1j})\partial_{x_j} + (c_{2j} - b_{1j})\partial_{y_j}, \quad J\partial_{x_k} = \partial_{y_k}$$

and so $N_J(\partial_{\varphi_1}, \partial_{x_j}) = \sum \alpha_j^k \partial_{x_k} + \beta_j^k \partial_{y_k}$ with constant coefficients α_j^k, β_j^k .

The last claim follows because $N_J(F, F) = 0$ and $\Pi = \mathbb{C}N_J(\partial_{\varphi_1}, F)$ is a linear subspace of F (of course invariant under J). Note that $\text{rk } \Pi$ can vary with $\varphi \in \mathcal{C}$. \square

Remark 3. *The conditions $N_J(F, F) = 0$ and $\text{Im } N_J \subset F$ allow to lift naturally this tensor from \mathcal{C} to E . Actually let $\xi \in T_x \mathcal{C}$ and $\eta, \theta \in T_x F \subset T_x E$. Let us choose some lift $\tilde{\xi} \in T_a E$ (i.e. $(\pi_*)\tilde{\xi} = \xi$). Let $\tilde{\eta}, \tilde{\theta} \in T_a F = F$ be identified with η, θ . Then we define $N_J(\tilde{\xi}, \tilde{\eta}) = \tilde{\theta}$ at $a \in E$ if $N_J(\xi, \eta) = \theta$ at $x = \pi(a)$. Our assumptions imply that this extension does not depend on a lift of ξ . So N_J on E is the canonical tensor.*

Consider now the case $\dim E = 4$. For every point $a \in E$ denote by $r = r_a \in T_a E$ the vertical vector equal to $x\bar{a} \in F \simeq T_a F$ with $x = \pi(a) \in \mathcal{C}$.

Theorem 9. *Let J be a linear bundle structure on a vector bundle $\pi : E \rightarrow \mathcal{C}$ with 2-dimensional fibers over a curve (\mathcal{C}, J') . Then for some complex structure J_0 on E , making the bundle holomorphic, we have:*

$$J = J_0 + \frac{1}{2}J_0N_J(r, \cdot).$$

Proof. Let us define the structure by the formula

$$J_0 = J - \frac{1}{2}JN_J(r, \cdot). \quad (7)$$

Since $N_J|_F \equiv 0$ this structure $J_0|_F = J|_F$ is linear on fibers. This proves the formula for J of the theorem.

We first show that the structure J_0 is almost complex. Note that by lemma 8 $N_J(r, \xi) \in F$ for any ξ and $N_J(r, \xi) = 0$ for $\xi \in F$. So

$$J_0^2 = J^2 - \frac{1}{2}J^2N_J(r, \cdot) - \frac{1}{2}JN_J(r, J\cdot) + \frac{1}{4}JN_J(r, JN_J(r, \cdot)) = J^2 = -\mathbf{1}.$$

Now we show that this J_0 is integrable. By Newlander-Nirenberg theorem [NW] this is equivalent to the vanishing of the tensor N_{J_0} . Let us choose local coordinates (z, φ) as in proposition 3. In these coordinates $J\partial_x = J_0\partial_x = \partial_y$ and we deduce from (7):

$$\begin{aligned} N_{J_0}(\partial_x, \partial_{\varphi_1}) &= N_J(\partial_x, \partial_{\varphi_1}) - [\partial_y, \frac{1}{2}JN_J(x\partial_x + y\partial_y, \partial_{\varphi_1})] \\ &\quad + J[\partial_x, \frac{1}{2}JN_J(x\partial_x + y\partial_y, \partial_{\varphi_1})] \\ &= N_J(\partial_x, \partial_{\varphi_1}) - \frac{1}{2}N_J(\partial_x, \partial_{\varphi_1}) - \frac{1}{2}N_J(\partial_x, \partial_{\varphi_1}) = 0. \end{aligned}$$

Since the bivector $\partial_x \wedge \partial_{\varphi_1}$ generates $\Lambda_{\mathbb{C}}^2 TE$ the claim follows. \square

Note that the tensor N_J on the right-hand side in the formula of the theorem can be defined only at the points of $\mathcal{C} \subset E$ and then lifted to E as in remark 3. Since one easily describes complex structures on vector bundles (for trivial bundles see §1.3) we get the complete description of linear bundle almost complex structures.

2.5 Normal form of 1-jet of an almost complex structure

Consider the ideal of real-valued functions corresponding to the curve \mathcal{C}

$$\mu = \{f \in C^\infty(M^4) \mid f(\mathcal{C}) = 0\}.$$

Degrees of this ideal give the filtration μ^k on every $C^\infty(M)$ -module, in particular we can talk about jets of tensor fields.

Theorem 10. *Let $\mathcal{C} \subset M^4$ be a pseudoholomorphic curve with respect to two almost complex structures J_1 and J_2 on M^4 . Assume $H^1(\mathcal{C}; N_{\mathcal{C}}M) = 0$, i.e. the curve is a disk D_R , the entire line \mathbb{C} or the sphere S^2 with trivial normal bundle $S^2 \cdot S^2 = 0$. If $J_1|_a = J_2|_a$ and $N_{J_1}|_a = N_{J_2}|_a$ at every point $a \in \mathcal{C}$ then the structures J_1 and J_2 are 1-equivalent, i.e. there exists a diffeomorphism ψ of a neighborhood $\mathcal{O}(\mathcal{C})$, preserving \mathcal{C} , such that*

$$J_2 = \psi^* J_1 \text{ mod } \mu^2.$$

Proof. This statement is an analog of theorem 1 [K1]. Details of the construction can be found there. Here we present a short proof with the indication of differences.

By the hypothesis the desired diffeomorphism ψ has 1-symbol $\Phi^{(1)} = \text{id} \in \tau^* \otimes \tau$ along the curve \mathcal{C} , where $\tau = TM|_{\mathcal{C}}$. Its 2-symbol $\Phi^{(2)} \in S^2 \tau^* \otimes \tau$ in local coordinates is given by $\Phi^{(2)}(\xi, \eta)^i = \frac{\partial^2 \psi^i}{\partial x^r \partial x^s} \xi^r \eta^s$. Moreover these symbols are compatible with the condition $\psi|_{\mathcal{C}} = \text{id}$:

$$\xi, \eta \in T\mathcal{C} \Rightarrow \Phi^{(2)}(\xi, \eta) = 0. \quad (8)$$

The symbol satisfies the following condition:

$$dJ_1(\xi, \eta) + J_1 \circ \Phi^{(2)}(\xi, \eta) = \Phi^{(2)}(J_2 \xi, \eta) + dJ_2(\xi, \eta),$$

where $d = d^\nabla$ is the differential w.r.t. some symmetric (and so not almost complex) curvature-free connection ∇ .

Equivalently this means that the tensor

$$P(\xi, \eta) = J_1 \circ \Phi^{(2)}(\xi, \eta) - \Phi^{(2)}(J_2 \xi, \eta) \quad (9)$$

satisfies the equation

$$P(\xi, \eta) = dJ_2(\xi, \eta) - dJ_1(\xi, \eta). \quad (10)$$

Now given J_1 and J_2 we have P and would like to find the corresponding Φ from (9). Form equality (10) we deduce

$$J_1 \circ P(\xi, \eta) = -P(J_2 \xi, \eta),$$

which implies $P(\xi, \eta) = J_1 B(\xi, \eta) - B(J_2 \xi, \eta)$ for some $(2, 1)$ -tensor B . Now the identity

$$P(\xi, \eta) - P(\eta, \xi) = P(J_2 \xi, J_2 \eta) - P(J_2 \eta, J_2 \xi) \quad (11)$$

implies the possibility to satisfy equation (9) by the choice

$$\Phi^{(2)}(\xi, \eta) = \frac{1}{2}[B(\xi, \eta) + B(\eta, \xi)] - \frac{J}{4}[B(\xi, J\eta) + B(\eta, J\xi) - B(J\xi, \eta) - B(J\eta, \xi)].$$

Identity (11) means exactly $N_{J_1} = N_{J_2}$. Note that the tensor B can be chosen to satisfy (8) and all the constructions respect this condition (TC is J_k -invariant). So we get $\Phi^{(2)}$ satisfying (8) and (9).

Now the symbols Φ stand for the differential of the mapping ψ we seek for. Calculations above show the $\tau^* \otimes \tau$ -valued 1-form generated by Φ is closed. Then the condition $H^1(\mathcal{C}; N_{\mathcal{C}}M) = 0$ imply that it is exact. Hence the symbols Φ are integrated to the diffeomorphism ψ we sought for. \square

Consider PH-sphere $\mathcal{C} = S^2 \subset M^4$, $\mathcal{C} \cdot \mathcal{C} = 0$. On the family of transversal disks D_φ , constructed in proposition 3, we can choose smooth global complex coordinate z so that z is complex w.r.t. the restriction $J|_{D_\varphi}$. So a neighborhood \mathcal{O} of S^2 is represented as a product $D^2 \times S^2$ with coordinates (z, φ) , φ taking values in \mathbb{C} . Let us define a complex structure by the product-formula $J_0 = J_v J_h$, with the vertical part J_v induced by the projection along the spheres $S^2 = \{z = \text{const}\}$ from the structure $J|_{D_\varphi}$ and the horizontal part J_h induced by the projection along the disks D_φ from the structure $J|_{\mathcal{C}}$.

Corollary 11. *Almost complex structure J in a neighborhood $\mathcal{O} \subset M^4$ of PH-sphere $\mathcal{C} = S^2$ is 1-equivalent to the almost complex structure $J' = J_0 + \frac{1}{2}J_0 N_J(r, \cdot) + \dots$, where N_J is the field of the Nijenhuis tensors along \mathcal{C} . In other words there exists a local diffeomorphism ψ , $\psi|_{\mathcal{C}} = \text{id}$, such that*

$$\psi^* J\xi = J_0\xi + \frac{1}{2}J_0 N_J(r_a, \xi) \text{ mod } \mu^2$$

for any vector $\xi \in T_a M$, $a \in \mathcal{O}$.

Proof. The values of the almost complex structure J and the almost complex mod μ^2 structure $\tilde{J} = J_0 + \frac{1}{2}J_0 N_J(r_a, \cdot) + \dots$ on the right-hand side of the formula are equal at the points of the sphere S^2 . Let us show that the same is true for their Nijenhuis tensors. Namely we show that the Nijenhuis tensor $N_{\tilde{J}}$ calculated according to definition (1) is equal to the prescribed Nijenhuis tensor N_J at all points S^2 .

The calculations are local and it is sufficient to consider the value of $N_{\tilde{J}}$ on two complex independent vectors, say on ∂_x and ∂_{φ_1} , where $\varphi = \varphi_1 + i\varphi_2$ and $z = x + iy$ are local coordinates defining J_0 . Let us denote by \doteq the equality mod μ . We have:

$$\begin{aligned} N_{\tilde{J}}(\partial_x, \partial_{\varphi_1}) &\doteq [J_0 \partial_x, \tilde{J} \partial_{\varphi_1}] - J_0[\partial_x, \tilde{J} \partial_{\varphi_1}] - J_0[J_0 \partial_x, \partial_{\varphi_1}] - [\partial_x, \partial_{\varphi_1}] \\ &\doteq [\partial_y, \partial_{\varphi_2} + \frac{1}{2}J_0 N_J(x\partial_x + y\partial_y, \partial_{\varphi_1})] \\ &\quad - J_0[\partial_x, \partial_{\varphi_2} + \frac{1}{2}J_0 N_J(x\partial_x + y\partial_y, \partial_{\varphi_1})] \\ &\doteq \frac{1}{2}J_0 N_J(\partial_y, \partial_{\varphi_1}) + \frac{1}{2}N_J(\partial_x, \partial_{\varphi_1}) \doteq N_J(\partial_x, \partial_{\varphi_1}). \end{aligned}$$

Thus $N_{\tilde{J}} = N_J$ along S^2 and the claim is proved. \square

Remark 4. Another way to construct a complex structure J'_0 in a neighborhood of PH-sphere $\mathcal{C} = S^2 \subset M^4$ with trivial normal bundle ($\mathcal{C} \cdot \mathcal{C} = 0$) is to foliate this neighborhood by homologous PH-spheres ([M2]) and to induce the vertical part J_v from some particular transversal disks D_{φ_0} .

Consider now arbitrary PH-curve \mathcal{C} with trivial normal bundle $\mathcal{C} \cdot \mathcal{C} = 0$. Similarly to the proposition 3 we find coordinates (z, φ) in a neighborhood of $\mathcal{C} \subset M$, where φ is multivalued and $\{\varphi = \text{const}\}$ is a family of transversal PH-disks D_φ . D_φ can be equipped with vector space structure and so for every $a \in D_\varphi \subset M^4$ we set $r \in T_a D_\varphi$ be the radius vector $\vec{0}z$ attached at the point a . Here we use the canonical identification $T_a F \simeq F$ for the vector spaces.

Theorem 9 expresses the linear bundle almost complex structure in terms of some complex structure and a Nijenhuis tensor. When almost complex structure is arbitrary then the conclusions of proposition 6 and theorem 1 are wrong (the formula of theorem 1 can give the operator with $J^2 \neq -1$). However on the level of 1-jet the theorem remains true.

Theorem 12. Let $\mathcal{C} \subset (M^4, J)$ be a PH-curve and $N_J \in \Lambda^2 \tau^* \otimes \tau$ be the field of Nijenhuis tensors of J along \mathcal{C} , where $\tau = TM|_{\mathcal{C}}$. Assume the Nijenhuis tensor characteristic distribution Π^2 is transversal everywhere to \mathcal{C} . Then for some complex structure J_0 in a neighborhood of \mathcal{C} we have:

$$J\xi = J_0\xi + \frac{1}{2}J_0N_J(r, \xi) \bmod \mu^2. \quad (12)$$

Proof. We can assume the disks D_φ have Π^2 as tangent planes at the points of \mathcal{C} . We use the symbol \doteq as in the proof of the corollary above. First note that formula (12) implies $J \doteq J_0$ for vertical vectors (belonging to TD_φ). Thus we define $J_0 \doteq J - \frac{1}{2}JN_J(r, \xi)$. Similarly to the proof of theorem 9 it is proved that $J_0^2 \doteq -1$ and that $N_{J_0} \doteq 0$. Thus J_0 is a complex structure $\bmod \mu^2$ [K1] and we can take instead of it any *complex* structure $\tilde{J}_0 = J_0 \bmod \mu^2$. \square

In this theorem J_0 is some complex structure while in the corollary 11 we had some definite complex structure. In the case $C = T^2$ we can also assume the complex structure J_0 is $J_{(\lambda)}$ from proposition ??.

Actually, if the complex structure J_0 in the torus neighborhood has normal bundle determined by the normal nonresonant pair (λ, ν) , then we can find coordinates (z, φ) in a neighborhood with the gluing rule (4) in which J_0 is standard. In these coordinates

$$J_\alpha^\gamma = \delta_\alpha^{\gamma+n} + \frac{1}{2}N_{\alpha\beta}^{\gamma+n}r^\beta + \bar{\delta}(|r|), \quad (13)$$

with $z = r^2 + ir^4$ and indices 1, 3 being used for the real and imaginary part of $\varphi = r^1 + ir^3$. Here $N_{\alpha\beta}^\gamma$ are the components of the Nijenhuis tensor. We assume every object changes its sign when one index is changed by 4, e.g. $N_{\alpha\beta}^{\gamma+4} = -N_{\alpha\beta}^\gamma$. The summation index β is assumed to take only values 2,4.

Remark 5. *Similar statement holds also for 1-jet of almost complex structure at a point. It says that for some coordinate system (x^i) near the point the operator J has components*

$$J_\alpha^\gamma = \delta_\alpha^{\gamma+n} + \frac{1}{4} N_{\alpha\beta}^{\gamma+n} x^\beta + \bar{\partial}(|x|), \quad (14)$$

where $N_{\alpha\beta}^\gamma$ are components of the Nijenhuis tensor and we assume $2n$ -twisted periodicity, i.e. $x^{i+2n} = -x^i$ etc.

This formula is obtained by studying the structural function (Weyl tensor) of the corresponding geometric structure ([K1],[KL]). The difference of the factors ($1/2$ instead of $1/4$) in the formulas is connected with the fact that we consider different coordinate systems: in (13) we consider the radial vector field r which connects the point to its projection on the torus and in (14) we use the usual radius-vector directing from a fixed origin. Such formulas for $J \bmod \mu^2$ can be considered along any PH-submanifolds of (M^{2n}, J) but generically there are only 0- and 2-dimensional PH-submanifolds.

3 Curves in a neighborhood of a PH-torus

3.1 Locally foliating families of cylinders

In this section we consider only pseudoholomorphic tori $\mathcal{C} = T^2$. Results of section 1 imply there are moduli of almost complex structures on a germ of \mathcal{C} contrary to the complex case in which Arnold found a normal form of a neighborhood of an elliptic curve $T^2 \subset M^4$. To get some analogs of the complex situation we consider foliations of a neighborhood of T^2 by pseudoholomorphic curves. Generically there are no compact curves in a neighborhood and PH-tori appear discretely because index of the corresponding linearized Cauchy-Riemann operator is zero ([Ku2]). So one should consider noncompact curves. We begin with PH-lines. Let $\|\cdot\|$ be some fixed norm.

Proposition 13. *There are two neighborhoods $\mathcal{O}' \subset \mathcal{O}$ of $T^2 \subset M^4$, a change of almost complex structure in $\mathcal{O} \setminus \mathcal{O}'$ and a number $C > 0$ with the following properties. For every $R > 0$ there exists a smooth family of PH-disks $f_\alpha : D_R \rightarrow \mathcal{O}$ with uniformly bounded norms $\|(f_\alpha)_*(z)\| \leq C$ and $\|(f_\alpha)_*(0)\| = 1$. Moreover this family fills some smaller neighborhood $\mathcal{O}'' \subset \mathcal{O}'$ of T^2 , i.e.*

$$\mathcal{O}'' \subset \cup_\alpha f_\alpha(D_R).$$

Proof. Let us take the universal covering $\hat{\mathcal{O}} \simeq \mathbb{C}D^2$ of \mathcal{O} . The torus is covered by the entire line $\mathbb{C} \rightarrow T^2$. Changing the structure J at infinity in $\hat{\mathcal{O}}$ and near the boundary to the integrable one we glue the manifold to the product $S^2 S^2$ with the line \mathbb{C} being glued to the first factor S_1^2 . Then the introduction of the taming symplectic product-structure $\omega = \omega_1 \oplus \omega_2$ yields a foliation of $S_1^2 S_2^2$ by PH-spheres S^2 in the homology class of the first factor if we additionally demand $\omega_1(S_1^2) < \omega_2(S_2^2)$, i.e. the homology class $[S_1^2]$ of the first sphere-factor

is symplectically simple. Here we use the fact that the dimension is 4: due to positivity of intersections [M1] we actually have a foliation ([M2]).

This foliation of S^2S^2 gives a family of big PH-disks on $\hat{\mathcal{O}}$ parametrized by the radius ρ of disk in \mathbb{C} out of which the almost complex structure is changed. To get estimates we use Brody reparametrization lemma as in [KO]. The filling property is given by the choice of perturbation of the structure in $\mathcal{O} \setminus \mathcal{O}'$: This is simple if we require the boundary of \mathcal{O} to be J -pseudoconvex ([G] 2.4.D). \square

We now consider filling by pseudoholomorphic cylinders $\mathcal{C}_R = [-R; R]S^1$. They can be considered also as annuli $\mathcal{C}_R \subset \mathbb{C} \setminus \{0\}$, but the next construction differs from the previous because the cylinders are now non-contractible in \mathcal{O} (Fig.1).

Proposition 14. *In the statement of proposition 13 we can change disks D_R to \mathcal{C}_R and get for every $R > 0$ a family of PH-cylinders $f_\alpha : \mathcal{C}_R \rightarrow \mathcal{O}$ with uniformly bounded norms and normalization $\|(f_\alpha)_*(0)\| = 1$. In addition this family is filling:*

$$\mathcal{O}'' \subset \cup_\alpha f_\alpha(\mathcal{C}_R).$$




Figure 1: Filling by PH-cylinders

Proof. Actually take a covering of the neighborhood \mathcal{O} which corresponds to one cycle of the torus. The torus is covered by the entire cylinder $\hat{\mathcal{C}} \rightarrow T^2$. We can change the almost complex structure J at infinity so that it makes possible to "pinch" each end of the cylinder. This means we perturb the structure J so that it is standard integrable outside some $\mathcal{C}_{R_2} \subset \hat{\mathcal{C}}$ and the support is also a big cylinder \mathcal{C}_{R_1} . Then we glue the ends to the disks. This operation gives us a sphere S^2 instead of the cylinder $\hat{\mathcal{C}} = \mathbb{R}S^1$. We can also assume that neighborhoods of two cylinder ends are pinched (Fig.2).

Thus we have a neighborhood U of the sphere S^2_0 . It is foliated by PH-spheres close to S^2_0 . Actually, we can change the structure J near the boundary of this neighborhood, glue and get the manifold-product $\hat{M} = S^2S^2$. As before it is foliated by PH-spheres. Thus U is foliated by PH-spheres and in the preimage they give a PH-foliation by cylinders. \square




Figure 2: Cutting and Gluing

Remark 6. Consider the infinite cylinder $\hat{C} = \mathcal{C}_\infty$. One can prove a more general statement: There exists a change of the structure J in $\mathcal{O} \setminus \mathcal{O}'$ such that \mathcal{O}' is filled with images of entire PH-cylinders $f_\alpha : \hat{C} \rightarrow \mathcal{O} : \mathcal{O}' \subset \cup_\alpha f_\alpha(\hat{C})$. The technical difficulty is however that we need reparametrization in the sequence of finite PH-cylinders passing the given point and the limiting curve can shift and not pass through the point (I am indebted to V.Bangert for a discussion about it). To achieve filling one needs to control the sequence. We will consider the problem elsewhere.

The positivity of intersections theory ([M1]) is applicable only to closed submanifolds and it cannot guarantee that leaves of the considered filling do not intersect. If there are no intersections of the leaves with close values of parameters and there are no self-intersections we call the family *locally foliating*.

Recall that PH-torus T^2 is parametrized by periodic coordinate $z \in \mathbb{C}$, i.e. the map $f_0 : \mathbb{C} \rightarrow T^2$ satisfies $f_0(z) = f_0(z + 2\pi) = f_0(z + \nu)$, where $2\pi, \nu$ are periods. We parametrize f_α by the condition $f_\alpha(z) = f_\alpha(z + 2\pi)$.

Fix some transversal PH-disk D_{φ_0} to the torus T^2 . Consider the return map $\Psi : D_{\varphi_0} \rightarrow D_{\varphi_0}$ for our family of surfaces f_α . This is defined as follows: if a point $z \in D_{\varphi_0}$ belongs to f_α , $z = f_\alpha(\varphi)$ we set $\Psi(z) = z'$ with $z' = f_\alpha(\varphi + \sigma(\varphi))$, $\sigma(\varphi) \approx \nu$, chosen to belong to D_{φ_0} again. This return map Ψ is determined non-uniquely since several cylinders of the family f_α can pass through z .

Proposition 15. Let the return map be either attracting or repelling, $|\Psi(z)/z| \in (0, \varepsilon)$ or $(\varepsilon^{-1}, \infty)$ for all $z \in D_{\varphi_0}$ close to zero and some fixed $\varepsilon < 1$ (we assume the inequality holds notwithstanding the nonuniqueness of Ψ). Then the constructed family $f_\alpha : \hat{C} \rightarrow \mathcal{O}$ is locally foliating in a neighborhood of T^2 .

Proof. The condition that Ψ is attracting or repelling means that there are no self-intersections of the leaves $f_\alpha(\hat{C})$. If a leaf $f_\alpha(\hat{C})$ transversally intersect another $f_\beta(\hat{C})$ with $\alpha \approx \beta$ then because of the convergence $f_\alpha(\mathcal{C}_R) \rightarrow f_\alpha(\hat{C})$ there is an intersection of the spheres $S^2 \subset \hat{M}$ from which cylinders are constructed. This is certainly impossible because our family of spheres is foliating. If the

leaves $f_\alpha(\hat{\mathcal{C}})$ and $f_\beta(\hat{\mathcal{C}})$ are tangent let us consider the first order of their jets which are different. This is the next number after the tangency order. Since the maps of finite cylinders \mathcal{C}_R tends to the maps of $\hat{\mathcal{C}}$ in C^∞ -topology the spheres in the manifold \hat{M} do intersect. Every such intersection even nontransversal contributes positively to the intersection number ([M1]) which contradicts $[S^2] \cdot [S^2] = 0$. It remains to note that if two pseudoholomorphic curves in an almost complex manifold have the infinite tangency they must coincide ([MS]). \square

For arbitrary almost complex structures it is not clear if there are foliations of the T^2 -neighborhood by cylinders. The following can be probably solved using a method similar to the Moser's proof [Mo] of PH-lines foliation persistence.

Question: Let T^2 be normal nonresonant elliptic curve. Let J be a small almost complex perturbation of the complex structure J_0 . The curve T^2 will be perturbed into close PH-curve \mathcal{C} . Is it true that a neighborhood $\mathcal{O}(\mathcal{C})$ is foliated by PH-cylinders?

3.2 Floquet theory for the PH-foliations

Holomorphic bundle (4) possesses a distinguished foliation $\{z = \text{const}\}$. We call this foliation the *foliation by λ -twisted cylinders*. Arnold's "Floquet-type" result in [A1]§27 implies that a neighborhood of every elliptic curve has a foliation by twisted cylinders which is biholomorphic to the standard one. Here we study PH-foliations.

Let $f_\alpha : \hat{\mathcal{B}} \rightarrow \mathcal{O}$ be a foliating family of a neighborhood of a PH-curve \mathcal{C} with trivial self-intersection. Let $D_\varphi = \{\varphi = \text{const}\}$ be a family of normal disks from proposition 3. Then every path $\gamma(t)$ on \mathcal{C} with $\gamma(0) = \varphi_0$, $\gamma(1) = \varphi_1$ gives a mapping $\Phi_\gamma : D_{\varphi_0} \rightarrow D_{\varphi_1}$ of shift along the leaves of f_α . For a loop γ we have an automorphism of D_φ . Since f_α is foliation there is no local holonomy: $\Phi_\gamma = \text{id}$ for contractible loops γ . Thus we can consider the map $\pi_1(\mathcal{C}) \rightarrow \text{Aut}(D_\varphi)$.

Definition 3. We call $\Phi_\gamma \in \text{Aut}(D_\varphi)$ the *monodromy map* along $\gamma \in \pi_1(\mathcal{C})$.

For example there is no monodromy for the sphere $\mathcal{C} = S^2$ and each choice of local coordinates in a normal disk D_{φ_0} gives coordinates for the others D_φ .

Let now $\mathcal{C} = T^2(2\pi, \nu)$. Since our foliating family f_α consists of cylinders there is no monodromy along one generating cycle, let along the cycle $\varphi \mapsto \varphi + 2\pi$. Denote by Φ_ν the monodromy along the other cycle $\varphi \mapsto \varphi + \nu$.

Let $f_\alpha : \hat{\mathcal{C}} \rightarrow \mathcal{O}$ be a foliating family of PH-cylinders in a neighborhood $\mathcal{O}' \subset \mathcal{O}$ of the PH-torus T^2 . Assume the monodromy of the family f_α is

$$\Phi_\nu(z) = \lambda z + \bar{\delta}(|z|). \quad (15)$$

Recall [A1] that $\lambda \in \mathbb{C} \setminus \{0\}$ is of (C, σ) -type if $|\lambda^m - 1| \geq C/m^{1+\sigma}$ for all $m \in \mathbb{Z}_{>1}$ (in particular such are all nonzero from Poincaré domain $|\lambda| \neq 1$).

Theorem 16. Let the monodromy Φ_ν be a germ of biholomorphic mapping with the number λ (15) of (C, σ) -type. Then the foliating family of cylinders f_α is similar to λ -twisted foliation in the following sense: There exist coordinates

(z, φ) with the gluing rule (4) such that transversal disks $D_\varphi = \{\varphi = \text{const}\}$ are pseudoholomorphic, φ is a complex coordinate on the curve \mathcal{C} , z is a complex coordinate on one D_{φ_0} and the foliation f_α is given by $\{z = \text{const}\}$.

Proof. A coordinate system z on a transversal disk D_{φ_0} provides coordinates on all others D_φ . These coordinates are multivalued because rotation along the second cycle $\varphi \mapsto \varphi + \nu$ yields the monodromy mapping $z \mapsto \psi(z) = \lambda z + \bar{\delta}(|z|)$.

Now we apply Poncaré-Siegel theorem ([A1]) in complex dimension 1 which states that the monodromy map is conjugate to the map $\psi(z) = \lambda z$. \square

3.3 Monodromy and transports

Unlike complex case almost complex monodromy can be non-holomorphic mapping of the fibers. Actually there are simple examples of PH-foliations with any prescribed monodromy.

Moreover even if the monodromy is complex (as required in theorem 16) the transport maps $\Phi_\gamma : (D_{\varphi_0}, J) \rightarrow (D_{\varphi_1}, J)$ can be not. In fact this is the occasion of $\text{codim} = \infty$.

Proposition 17. *Let $\mathcal{C} \subset M$ be a PH-curve in a 4-dimensional manifold and let $f_\alpha : P \rightarrow \mathcal{O}$ be a local PH-foliating family of some neighborhood $\mathcal{O}(\mathcal{C})$. Let also D_φ be a transversal PH-foliation and Φ_γ be the corresponding transport map $D_{\varphi_0} \rightarrow D_{\varphi_1}$ along curves $\gamma(t) \subset \mathcal{C}$, $\gamma(0) = \varphi_0$, $\gamma(1) = \varphi_1$. If all Φ_γ are holomorphic then the Nijenhuis tensor characteristic distribution Π^2 is tangent to the leaves of f_α .*

Proof. Actually the foliation provides a local bundle $\pi : \mathcal{O} \rightarrow D_\varphi$. The hypothesis that all transports are complex is equivalent to the claim that the bundle π with fibers $f_\alpha(P)$ is almost complex. Therefore the statement follows from proposition 6. \square

Remark 7. *If the distribution Π^2 is not integrable the transports are not complex. But even integrability is not sufficient, see §2.2.*

3.4 PH-tori deformation problem

Generically holomorphic 2-torus in a complex manifold cannot be deformed ([A1]). The same situation is also with PH-tori in almost complex manifolds. This follows from vanishing of the index of the linearized Cauchy-Riemann operator ([Ku1]). By the deformation we mean existence of close homologous PH-torus of the same periods $T^2 = T^2(2\pi, \nu)$. In this section we consider some examples where we can make the condition of non-existence explicit.

$\underline{1}^\circ$) Let us consider linear bundle almost complex structure J on the bundle $E \rightarrow T^2$. There exist coordinates (z, φ) with the gluing rule (4) such that

$$\begin{cases} J\partial_x = \partial_y, & J\partial_{\varphi_1} = \partial_{\varphi_2} + x \cdot v - y \cdot Jv, \\ J\partial_y = -\partial_x, & J\partial_{\varphi_2} = -\partial_{\varphi_1} - x \cdot Jv - y \cdot v. \end{cases}$$

This formula follows from theorem 9 and the coordinates are determined by J_0 . Vector field $v = \frac{1}{2}JN_J(\partial_x, \partial_{\varphi_1})$ can be decomposed $v = \alpha\partial_x + \beta\partial_y$ with $\alpha = \alpha(\varphi)$, $\beta = \beta(\varphi)$. The complexified vector bundle is decomposed $T_{\mathbb{C}}E = E_+ + E_-$, where $E_{\pm} = \{\xi \mid J\xi = \pm i\xi\}$; $E_- = \bar{E}_+$. Vectors

$$U_1 = \partial_{\varphi} - z\bar{b}\partial_{\bar{z}}, \quad U_2 = \partial_z,$$

form a basis of E_+ . Here $\partial_{\varphi} = \frac{1}{2}(\partial_{\varphi_1} - i\partial_{\varphi_2})$, $\partial_z = \frac{1}{2}(\partial_x - i\partial_y)$ and $\bar{b} = \frac{\beta + i\alpha}{2}$. Thus the basis of E_+^* in the decomposition $T_{\mathbb{C}}^*E = E_+^* + E_-^*$ ($E_-^* = \bar{E}_+^*$) is

$$\omega_1 = d\varphi, \quad \omega_2 = dz + \bar{z}b d\bar{\varphi}.$$

Now every pseudoholomorphic torus in E homologous to the zero section T^2 is of the form $f(T^2)$ for some section f of the bundle $E \rightarrow T^2$. This is to say each PH-torus in E has unique transversal intersection with every fiber. Actually we may compactify the fibers of the bundle to the spheres and the claim follows from the positivity of intersections (or even simpler by studying the degree of the projection of this torus to the torus-base).

Let us deduce the equation for f . The curve $f(T^2)$ is pseudoholomorphic iff

$$\omega_2|_{z=f(\varphi)} = c \cdot \omega_1.$$

Substituting $df = f_{\varphi}d\varphi + f_{\bar{\varphi}}d\bar{\varphi}$ we have:

$$f_{\bar{\varphi}} + b\bar{f} = 0. \tag{16}$$

Theorem 18. *Let J be a linear bundle almost complex structure and J_0 be the corresponding complex structure from the decomposition of theorem 9. Suppose the number λ , determined by the complex structure J_0 in the bundle E via (4), is of unit length: $|\lambda| = 1$. Assume also that the function $\Lambda \in C^{\infty}(T^2)$, determined uniquely by the equation $\frac{1}{2}JN_J(\partial_z, \partial_{\varphi}) = \Lambda\partial_{\bar{z}}$, is nonzero holomorphic function: $\partial_{\bar{\varphi}}\Lambda = 0$, $\Lambda \neq 0$. Then zero section T^2 is the unique PH-torus in E .*

Proof. First note that since

$$\begin{aligned} \frac{1}{2}JN_J(\partial_z, \partial_{\varphi}) &= (\alpha - i\beta)\partial_{\bar{z}}, & \frac{1}{2}JN_J(\partial_{\bar{z}}, \partial_{\varphi}) &= 0, \\ \frac{1}{2}JN_J(\partial_z, \partial_{\bar{\varphi}}) &= 0, & \frac{1}{2}JN_J(\partial_{\bar{z}}, \partial_{\bar{\varphi}}) &= (\alpha + i\beta)\partial_z, \end{aligned} \tag{17}$$

we have $\Lambda = -2i\bar{b}$. Thus $\Lambda_{\bar{\varphi}} = 0 \Leftrightarrow \bar{\Lambda}_{\varphi} = 0 \Leftrightarrow b_{\varphi} = 0$.

Let us show the equation (16) has no nonzero solutions. Complex Laplacian of f equals $f_{\bar{\varphi}\varphi} = -b\bar{f}_{\varphi} = -b\bar{f}_{\bar{\varphi}} = |b|^2f$. Our torus neighborhood is the trivial cylinder $\mathcal{C}^2 = \{\varphi \in \mathbb{C} \mid \text{Im } \varphi \in [0, \text{Im } \nu)\} / 2\pi\mathbb{Z}$ neighborhood glued by the rule $(z, \varphi) \mapsto (\lambda z, \varphi + \nu)$. Thus when $\varphi \mapsto \varphi + \nu$ we have: $f \mapsto \lambda f$. So integrating over the cylinder gives:

$$\begin{aligned} \int_{\mathcal{C}^2} (f_{\bar{\varphi}\varphi}\bar{f} + f_{\varphi}\bar{f}_{\bar{\varphi}}) \frac{i}{2}d\varphi \wedge d\bar{\varphi} &= \int_{\mathcal{C}^2} \frac{\partial}{\partial\varphi}(f_{\bar{\varphi}}\bar{f}) \frac{i}{2}d\varphi \wedge d\bar{\varphi} = \int_{\mathcal{C}^2} \frac{i}{2}d(f_{\bar{\varphi}}\bar{f}d\bar{\varphi}) = \\ &= (\lambda\bar{\lambda} - 1) \oint_{S^1} \frac{i}{2}f_{\bar{\varphi}}\bar{f}d\bar{\varphi} = 0. \end{aligned}$$

So using the calculation with the Laplacian we have:

$$\int_{\mathcal{C}^2} (|b|^2|f|^2 + |f_{\bar{\varphi}}|^2) d\varphi_1 \wedge d\varphi_2 = 0. \quad (18)$$

Therefore since $|b| \neq 0$ we get $f = 0$. Thus there are no homologous to the zero section PH-tori \tilde{T}^2 with $f \neq 0$. If the homology class of \tilde{T}^2 is a multiple of the zero section $[\tilde{T}^2] = k[T^2]$ a k -finite covering finishes the proof. \square

Remark 8. *If $|b| = 0$, i.e. almost complex structure J is integrable $J = J_0$, equality (18) implies that f is holomorphic section. Thus if $\lambda^n \neq 1$ we get again $f = 0$ comparing the Fourier coefficients of f .*

2°) Consider a general almost complex structure J with Nijenhuis tensor characteristic distribution Π^2 transversal to some PH-torus T^2 . The linearized equation for close PH-tori can be written in the form

$$f_{\bar{\varphi}} + af + b\bar{f} = 0. \quad (19)$$

Actually, the linearization does not depend on a change of the structure J by second order quantities. Thus we can perturb J to make the distribution Π^2 integrable in $\mathcal{O} \supset T^2$. This new almost complex structure is given by the formula (5).

Let's write the equation for close PH-tori. The basis of E_+ is

$$U_1 = \partial_{\varphi} + \frac{\bar{A}}{A+2i} \partial_{\bar{\varphi}} + \frac{\bar{B}}{A+2i} \partial_{\bar{z}}, \quad U_2 = \partial_z,$$

where $A = A_1 + iA_2$, $B = B_1 + iB_2$. The corresponding basis of E_+^* is

$$\omega_1 = d\varphi - \frac{A}{\bar{A}-2i} d\bar{\varphi}, \quad \omega_2 = dz + \frac{1}{4} \frac{\bar{A}B}{A_2+1} d\varphi - \frac{1}{4} \frac{(A+2i)B}{A_2+1} d\bar{\varphi}.$$

So the equation $(\omega_2 - c \cdot \omega_1)|_{z=f(\varphi)} = 0$ implies the required equation on f :

$$f_{\bar{\varphi}} + \frac{A}{\bar{A}-2i} f_{\varphi} - \frac{B}{\bar{A}-2i} = 0. \quad (20)$$

Denote by A^0 and B^0 linearizations by fiber coordinate of the functions A and B respectively. Note that linearization of equation (6) implies that A^0 is holomorphic w.r.t. z , that is we can bring our equations to the constant A^0 .

Since $A|_{z=0} = 0$, $B|_{z=0} = 0$, linearization of equation (20) is

$$f_{\bar{\varphi}} - \frac{i}{2} B^0(f) = 0,$$

which has the form (19) if we set $-\frac{i}{2} B^0 = az + b\bar{z}$.

Since we have equations (6) the Nijenhuis tensor of (5) is

$$N_J(\partial_x, \partial_{\varphi_1}) = \left(\frac{\partial B_1}{\partial y} - \frac{\partial A_1}{\partial x} B_1 + \frac{\partial A_2}{\partial x} \frac{A_1 B_1 - B_2}{A_2 + 1} + \frac{\partial B_2}{\partial x} \right) \partial_x \\ + \left(\frac{\partial B_2}{\partial y} - \frac{\partial A_1}{\partial x} B_2 + \frac{\partial A_2}{\partial x} \frac{A_1 B_2 + B_1}{A_2 + 1} - \frac{\partial B_1}{\partial x} \right) \partial_y.$$

Therefore linearizing A and B we conclude that its values on T^2 are given by the formula (see also (17))

$$\frac{1}{2} J N_J(\partial_z, \partial_{\varphi}) = -2i\bar{b} \partial_{\bar{z}}.$$

Remark 9. *Since the only invariant of 1-jet of J on a PH-curve is the Nijenhuis tensor, which we expressed by the function $b(\varphi)$, we can bring the function $a(\varphi)$ in (19) to the simplest form. Namely we can introduce coordinates (φ, z) using the complex structure J_0 of theorem 12. This gives $a = 0$ for normal coordinate z on the torus with gluing rule (4). Alternatively we can have global well-defined coordinate z but then $a = \text{const}$. This proves a suggestion on p. 430 [Mo] that "the linearized equation can be brought into the form (19) with $a = \text{const}$ ".*

Theorem 19. *Let almost complex structure J in a neighborhood of PH-curve T^2 be described by formula (12) with complex structure J_0 having $|\lambda| = 1$. If the characteristic distribution Π^2 is transversal to T^2 and for linearized structure $b(\varphi)$ is anti-holomorphic, then the curve T^2 is isolated and persistent.*

Proof. Actually as Moser [Mo] noticed if the linearized equation $f_{\bar{\varphi}} + af + b\bar{f} = g$ has a unique solution for any $g \in C^\infty(T^2; \mathbb{C})$ then the torus is isolated and persistent. But the linearization we studied in theorem 18. \square

3° Note that in holomorphic bundle with $\lambda^n = 1$ one can find tori $f(T^2)$ with $f \neq 0$ of the type $T^2(2\pi kn, \nu l)$, which cover zero section torus $T^2 = T^2(2\pi, \nu)$. But in a fixed homology class all PH-tori are of the same holomorphic type:

Lemma 20. *Let $\tilde{T}_1^2, \tilde{T}_2^2 \subset E$ be two PH-tori in a linear almost complex bundle $E \rightarrow T^2$. If they are homologous then they are biholomorphic.*

Proof. First consider tori in the homology class of the zero section T^2 . As was shown before theorem 18 the projection is a diffeomorphism. Since in linear almost complex bundles $E \rightarrow T^2$ the projection is an almost complex mapping, its restriction is a biholomorphism of the tori: $\tilde{T}_1^2 \simeq T^2 \simeq \tilde{T}_2^2$. The general case $[\tilde{T}_i^2] = k[T^2]$ follows from the case $k = 1$ by means of a k -covering. \square

If we do not demand the bundle condition the opposite situation can occur: in example [A1]§27 a neighborhood of the torus is foliated by holomorphic tori of different holomorphic type. Similar situation occurs also in almost complex case and invariants of section 1 can be nontrivial:

Example. Consider a foliation $f_\alpha : T^2 \rightarrow \mathcal{O}$ of 4-dim neighborhood of some torus. Introduce the structure J in horizontal directions so that all the tori T_α^2 are pseudoholomorphic but nonequivalent (the parameter ν is changing). Choose the structure J on the normals D_φ so that the transports Φ_γ are not holomorphic (§3.3). Define J globally by the product formula. Then the distribution Π^2 is nonintegrable and we get the distribution $L^1 = \Pi^3 \cap T(T^2)$ (possibly with singularities).

4°) Note that the tori in the same homology class can occur both in families and discretely.

Example. Let (T^4, J_0) be the standard complex torus, i.e. quotient of \mathbb{C}^2 by the lattice \mathbb{Z}^4 . Consider a PH-torus $T_0^2 \subset T^4$. It is possible to perturb J_0 in a neighborhood \mathcal{O} of T_0^2 so that the new structure J is isomorphic to the model structure near the torus $T^2(2\pi, \nu)$ given by (4) in a neighborhood $\mathcal{O}' \subset \mathcal{O}$ and $J = J_0$ outside \mathcal{O} . Then there is 2-parametric family of PH-tori of $[T_0^2]$ -homology class outside \mathcal{O} and a unique PH-torus T_0^2 inside \mathcal{O}' .

5°) Note that for PH-torus $\mathcal{C} = T^2 \subset (M^4, J)$ with $N_J|_a = 0$ for all $a \in T^2$ the normal bundle $N_{\mathcal{C}}M$ is holomorphic.

Proposition 21. *If the Nijenhuis tensor vanishes along a PH-torus and the pair (λ, ν) , characterizing the holomorphic bundle $N_{\mathcal{C}}M$, is nonresonant then small neighborhood \mathcal{O} of this torus cannot be foliated by $T^2(2\pi, \nu)$ -tori.*

Proof. Actually if there is a PH-foliation by tori then the linearization of this foliation determines a holomorphic foliation of the normal bundle which is impossible by [A1]§27. \square

6°) An intermediate condition on foliation between holomorphic and pseudoholomorphic is that it be pseudoholomorphic with complex transports. Proposition 17 implies

Proposition 22. *If the distribution Π^2 in a neighborhood $\mathcal{O} \supset T^2$ is not integrable or is integrable with noncompact leaves, then \mathcal{O} cannot be foliated by PH-tori with complex transports.*

Note that in the considerations above we needed to fix a holomorphic structures on the tori sought for. The last proposition does not require this.

A Normal bundle in Riemannian geometry.

The construction of structure on the normal to the submanifold bundle from §2.1 can be carried out for some other cases.

For example in Riemannian geometry there exists a unique Riemannian connection ∇ . This Levi-Civita connection splits the normal bundle $N_L M$ of any submanifold $i : L \subset M$ with induced Riemannian metric $g_L = i^*(g_M)$. However

the uniqueness of the connection ∇ makes the situation more rigid and for our construction we should require that L is a totally geodesic submanifold.

So we construct the metric in $N_L M$ and then again ask about relations between two tensors: Riemannian curvature R_g of the manifold M at points of $L \subset M$ and the curvature \hat{R}_g for the total space of $N_L M$ at zero section $L \subset N_L M$. In general there are no relations, but for some parts of the tensors there is.

Namely consider the curvature of the normal bundle R^\perp , which is the curvature tensor of the normal connection ∇^\perp given by the orthogonal decomposition in $TM|_L = TL \oplus N_L M$, $W = W_\parallel + W_\perp$. Note that $R^\perp(X, Y) = \hat{R}_g(X, Y)$ for $X, Y \in TL$ and the left-hand side is not defined for others X, Y .

Let $\Pi : TL \otimes TL \rightarrow N_L M$ be the second quadratic form of L and $A : TL \otimes N_L M \rightarrow TL$ be the shape (Peterson) operator given by $g(A(X, V), Y) = g(\Pi(X, Y), V)$, $X, Y \in TL$, $V \in N_L M$. Then if we denote R^L the curvature of the Levi-Civita connection of L we have the following famous equations:

$$\begin{aligned} [R_g(X, Y)Z]_\parallel &= R^L(X, Y)Z + A(Y, \Pi(X, Z)) - A(X, \Pi(Y, Z)) && \text{(Gauss eq.)}, \\ [R_g(X, Y)Z]_\perp &= (\nabla_Y \Pi)(X, Z) - (\nabla_X \Pi)(Y, Z) && \text{(Codazzi-Maynardi eq.)}, \\ [R_g(X, Y)V]_\perp &= R^\perp(X, Y)V + \Pi(X, A(Y, V)) - \Pi(Y, A(X, V)) && \text{(Ricci eq.)}, \end{aligned}$$

where $X, Y, Z \in TL$, $V \in N_L M$.

In particular when L is totally geodesic $\Pi = 0$ and $A = 0$, so that the equations above mean $R_g(X, Y) = \hat{R}(X, Y)$ for $X, Y \in TL$ at the points of L . However there are no relations if we allow X, Y to be arbitrary from TM .

So horizontal parts of the both curvatures R_g and \hat{R}_g coincide. Note that in the almost complex case this is trivially so because these horizontal parts vanish.

Question: What happens in other geometries – conformal, projective etc? There are also notions of normal connections ($[N]$) but on the normal spaces considered as bundles not manifolds.

Acknowledgements. I would like to thank V.I. Arnold who addressed me with several questions and suggestions originating the paper. During the work I had the benefit of discussions Riemannian geometry with D.V. Alekseevskij and projective geometry with V. Goldberg. I also express my thanks to V. Bangert for inviting me to the University of Freiburg and for valuable discussions during the visit.

Note on figures: if they are not visible look at www.math.uit.no/seminar/preprints.html

References

- [A1] V.I. Arnold, "Additional chapters of the theory of ordinary differential equations", Moscow, Nauka (1978) [Russian]; Engl. transl. "Geometrical methods in the theory of ordinary differential equations", Springer-Verlag, New York-Berlin, (1988).

- [A2] "Arnold's Problems", Moscow, Phasis (2000) [Russian; English transl. to appear].
- [B] V. Bangert, "Existence of a complex line in tame almost complex tori", *Duke Math. J.*, **94**, no.1 (1998), 29–40
- [BCG] R. L. Bryant, S. S. Chern, R. B. Gardner, H. L. Goldschmidt, P. A. Griffiths, "Exterior differential systems", Mathematical Sciences Research Institute Publications, **18**, Springer-Verlag, New York, (1991).
- [G] M. Gromov, "Pseudo-holomorphic curves in symplectic manifolds", *Invent. Math.*, **82** (1985), 307–347
- [KL] B. S. Kruglikov, V. V. Lychagin, "On equivalence of differential equations" *Acta et Commentationes Universitatis Tartuensis de Mathematica*, v.3 (1999), 7–29
- [KO] B. Kruglikov, M. Overholt, "The Kobayashi pseudodistance on almost complex manifolds", *Differential Geom. Appl.* v.11 (1999), 265–277
- [K1] B. S. Kruglikov, "Nijenhuis tensors and obstructions for pseudoholomorphic mapping constructions", *Mathematical Notes*, **63**, no.4 (1998), 541–561
- [K2] B. S. Kruglikov, "Some classificational problems in four dimensional geometry: distributions, almost complex structures and Monge-Ampere equations", *Math. Sbornik*, **189**, no.11 (1998), 61–74
- [K3] B. S. Kruglikov, "Existence of close pseudoholomorphic disks for almost complex manifolds and application to the Kobayashi-Royden pseudonorm", *Functional Analysis and Appl.* **33**, no.1 (1999), 46–58
- [Ku1] S. B. Kuksin, "Pseudoholomorphic 2-tori in T^4 ", *Differential Geom. Appl.*, **6**, no. 2 (1996), 109–119
- [Ku2] S. B. Kuksin, "Double-periodic solutions of nonlinear Cauchy-Riemann equations", *Comm. Pure Appl. Math.*, XLIX (1996), 639–676
- [L] A. Lichnerowicz, "Théorie globale des connexions et des groupes d'holonomie", Roma, Edizioni Cremonese (1955)
- [M1] D. McDuff "The local behavior of holomorphic curves in almost complex 4-manifolds", *Journ. Diff. Geom.*, **34** (1991), 143–164
- [M2] D. McDuff "Blow ups and symplectic embeddings in dimension 4", *Topology*, **30**, no. 3 (1991), 409–421
- [MS] D. McDuff, D. Salamon, "J-holomorphic curves and Quantum cohomology", Providence, RI, AMS, *University Lecture Series*, **6** (1994)

- [Mo] J. Moser, "*On the persistence of pseudoholomorphic curves on an almost complex torus*", *Invent. Math.*, **119** (1995), 401–442
- [N] A. P. Norden, "*Spaces with affine connection*" (Russian), "Nauka", Moscow (1976).
- [NW] A. Nijenhuis, W. Woolf, "*Some integration problems in almost-complex and complex manifolds*", *Ann. Math.*, **77** (1963), 424–489
- [S] S. Sternberg, "*Lectures on Differential geometry*", Prentice-Hall, New Jersey (1964).

Department of Math. and Stat., University of Tromsø, Norway
Mathem. Modelling Chair, Moscow Baumann State Technological University, Russia
E-mail: kruglikov@math.uit.no