Invariants of geometric structures on differential systems (distributions)

by

Thomas Opsahl

Cand. Mag. , University of Tromso (1999)

Submitted to the Department of Mathematics
in partial fulfillment of the requirements for the degree of

Master of Science

at the

UNIVERSITY OF TROMSO

June 2002

© Remove to use default copyright from the style

The author hereby grants to University of Tromso permission to reproduce and
to distribute copies of this thesis document in whole or in part.

Signature of Author .....
Department of Mathematics
21 June 2002

Certified by.....
David Collis
Director, Sound and Furry Laboratory
Research Head

Certified by.....
Boris Kruglikov
Associate Professor
Thesis Supervisor

Accepted by .....
Arthur C. Clarke
Chairperson, Department Committee on Graduate Students

Contents

1 Introduction **3**

2 Preliminaries **5**
2.1 Symplectic Manifolds 5
2.2 Contact Manifolds 8
2.3 Jet bundles π_k 9
2.4 Monge-Ampere equations and effective forms 10
2.5 Classification of effective forms 14

3 Hyperbolic M-A equations and 5-dimensional strict contact manifolds **17**
3.1 The extending sequence of derived submodules 17
3.2 Two canonical decompositions of τ_M 18
3.3 The Nijenhuis tensor 24

4 Invariants of hyperbolic M-A equations **31**
4.1 The canonical 1-form σ and the distribution Γ^1_σ 31
4.2 The vectorvalued 2-form R^σ_j 34
4.3 The canonical $\{e\}$-structure 40

5 Appendix A **44**

6 Appendix B **47**

Chapter 1

Introduction

In this thesis we deal with classification of second order Monge-Ampere equations defined on a 2-dimensional smooth manifold M. These are partial differential equations of the form

$$F_1 + F_2\frac{\partial^2 u}{\partial q_1^2} + F_3\frac{\partial^2 u}{\partial q_2^2} + F_4\frac{\partial^2 u}{\partial q_1 \partial q_2} + F_5\left(\frac{\partial^2 u}{\partial q_1^2}\frac{\partial^2 u}{\partial q_2^2} - \left(\frac{\partial^2 u}{\partial q_1 \partial q_2}\right)^2\right) = 0, \qquad (1.1)$$

where the coefficients are functions in $q_1, q_2, u(q_1, q_2), \frac{\partial u}{\partial q_1}, \frac{\partial u}{\partial q_2}$ and q_1, q_2 are the coordinate functions on M. Monge-Ampere equations (with arbitrary number of variables) have occupied both physicists and mathematicians for several centuries. Some of the earliest results belong to Goursat ([G]) and Lie ([Lie]) at the end of the 19th century. Our investigation takes place in a geometrical framework with roots going back to Sophus Lie. The geometrical approach was proposed by V. Lychagin in [Ly1]. It gives us a way to represent the equations as geometrical structures, so called effective forms, on the manifold of 1-jets J^1M.

In chapter 2 we briefly recall the basic definitions from symplectic and contact geometry. Specially we introduce the strict contact manifold J^1M. The contact form on J^1M we denote by ω. This contact form induce a contact distribution $\mathcal{C}$ on J^1M. The last part of the chapter concerns with the geometrical representation of equation 1.1 in the strict contact manifold J^1M. We show that a given Monge-Ampere equation can be represented as a so called conformal effective 2-form $\theta \in \Lambda^2\mathcal{C}^*$ ([Ly1], [LRC]). We also give the pointwise classification of effective forms described in [LRC]. The chapter ends with the definition of generalized Monge-Ampere equations as a pair (ω, θ).

In chapter 3 we begin our investigation of a 5-dimensional strict contact manifold (M, ω) together with a conformal effective form θ. The rank 4 contact distribution induced from ω is denoted by Π. We start the chapter by recalling that any distribution induces the so called extending sequence of derived submodules. Next we introduce the operator $j : \Pi \longrightarrow \Pi$ induced by ω and θ. This operator splits the distribution

Π into two distributions Π^2_+ and Π^2_- of rank 2. We show that there are two canonical decompositions of the tangent bundle τ_M. In the last section of the chapter we define the Nijenhuis tensor $N_j \in \Lambda^2\Pi^* \otimes \Pi$ and we investigate its properties.

In chapter 4 we introduce the two invariants σ, R^σ_j of the generalized Monge-Ampere equation. We investigate how σ and R^σ_j relate to the Nijenhuis tensor and the distributions from the two decompositions of τ_M. The chapter ends with a classifying theorem where we state that nondegenerate, generalized Monge-Ampere equations defines a canonical frame on M.

Chapter 2

Preliminaries

In this chapter we establish a correspondence between Monge-Ampere equations and so called effective forms. We also introduce notions that we need in the main part of this thesis. In what follows we consider the $\mathbb{R}$-algebra $C^\infty(M)$. The module of sections of a bundle π over M is denoted by $C^\infty(\pi)$.

2.1 Symplectic Manifolds

Let V be a vector space. Then we call $\mu \in \Lambda^2 V^*$ **nondegenerate** if for any vector $X \in V$ there exists some vector $Y \in V$ such that $\mu(X, Y) \neq 0$.

Definition 2.1 *Let V be a vector space and let $\Omega \in \Lambda^2 V^*$ be nondegenerate. Then we call the pair (V, Ω) a **symplectic vector space** and we refer to Ω as the **symplectic form**.*

Definition 2.2 *Let (V, Ω) be a symplectic vector space and let W be a subspace in V. We say that W is a **Lagrangian subspace** if W is skew orthogonal to itself with respect to the symplectic form Ω. That is:*

$$W^{\perp_\Omega} = \{ v \in V | \, \Omega(v, w) = 0 \ \forall \ w \in W \} = W.$$

As a consequence of the definition all symplectic vector spaces have to be even dimensional. Consider now a smooth manifold M. It is well known that the space of smooth functions over M, denoted $C^\infty(M)$, is an $\mathbb{R}$-algebra. For each point x in M, the space $\mu_x = \{ f \in C^\infty(M) | \, f(x) = 0 \}$ is an ideal in $C^\infty(M)$. This ideal defines the following filtration of ideals:

$$C^\infty(M) \supset \mu_x \supset \mu_x^2 \supset \mu_x^3 \supset \cdots .$$

The quotientspace

$$T_x^* M = \frac{\mu_x}{\mu_x^2}$$

is the cotangent space of M at the point x. T_x^*M is a vector space with the dimension equal to the dimension of M. The dual vector space T_xM is the tangent space of M at the point x.

Definition 2.3 *Let* $TM = \bigcup_{x\in M} T_xM$ *and* $T^*M = \bigcup_{x\in M} T_x^*M$. *We call the dual vector bundles*

$$\tau_M : TM \longrightarrow M$$

and

$$\tau_M^* : T^*M \longrightarrow M$$

the ***tangent*** *and* ***cotangent bundle*** *to* M *correspondingly.*

The $C^\infty(M)$-module of sections of the vector bundle $\Lambda^i\tau_M^*$ we denote by $\Omega^i(M)$, while the module of sections of the dual bundle $\Lambda^i\tau_M$ we denote by $\mathcal{D}_i(M)$.

Definition 2.4 *Let* M *be a* $2n$*-dimensional manifold and let* $\Omega \in \Omega^2(M)$ *be nondegenerate, i.e.* $\Omega_x \in \Lambda^2 T_x^*M$ *is nondegenerate for all* x *in* M. *Then we say that* (M, Ω) *is a* ***symplectic manifold****.*

Example 2.5 *Let* $M = \mathbb{R}^{2n}(q_1, \ldots, q_n, p_1, \ldots, p_n)$. *Then*

$$T_x^*M \simeq \mathbb{R}^{2n}(d_xq_1, \ldots, d_xq_n, d_xp_1, \ldots, d_xp_n)$$

and one can check that

$$\Omega_x = d_xq_1 \wedge d_xp_1 + \cdots + d_xq_n \wedge d_xp_n$$

defines a nondegenerate 2-form $\Omega \in \Omega^2(M)$*, such that* (M, Ω) *is a symplectic manifold.*

Definition 2.6 *The* ***vector bundle*** $\pi : E \longrightarrow B$ *is called* ***symplectic*** *if it is equipped with a symplectic structure* Ω_x *in fibers* $\pi^{-1}(x)$ *smoothly depending on* x.

Definition 2.7 *Let* (π, Ω) *be a symplectic vector bundle and let* γ *be a subbundle of* π. *We say that* γ *is a* ***Lagrangian subbundle*** *of* π *if every fiber in* γ *is a Lagrangian subspace of the corresponding fiber in* π.

A nondegenerate form $\Omega \in \Omega^2(M)$ defines the isomorphism

$$\begin{array}{rccccc} \Gamma: & \mathcal{D}_1(M) & \longrightarrow & \Omega^1(M) & \\ & \xi & \longmapsto & i_\xi\Omega & = \Omega(\xi, \bullet) \end{array}$$

of $C^\infty(M)$-modules.

Definition 2.8 *Let*

$$\pi_1 : E_1 \longrightarrow B_1 \qquad \text{and} \qquad \pi_2 : E_2 \longrightarrow B_2$$

be vectorbundles and let us denote fibers by F_1, F_2 in E_1, E_2 respectively. By ***morphism of vectorbundles*** *$\pi_1 \longrightarrow \pi_2$, we mean a pair $\left(h, \widehat{h}\right)$ of smooth maps such that the diagram*

$$\begin{array}{ccc} E_1 & \xrightarrow{\hat{h}} & E_2 \\ {\scriptstyle \pi_1}\uparrow & & \downarrow{\scriptstyle \pi_2} \\ B_1 & \xrightarrow{h} & B_2 \end{array}$$

commutes and

$$\widehat{h} : F_1(x) \longrightarrow F_2(h(x))$$

is a linear map for all points x in B_1.

When h is a diffeomorphism and $\widehat{h} : F_1(x) \longrightarrow F_2(h(x))$ is a linear isomorphism for all x in B_1, we say that $\left(h, \widehat{h}\right)$ is an isomorphism of vectorbundles. The following lemma is well known.

Lemma 2.9 *Let π_1, π_2 be the vectorbundles.*

$$\pi_1 : E_1 \longrightarrow B, \qquad \pi_2 : E_2 \longrightarrow B.$$

Let

$$\varphi : C^\infty(\pi_1) \longrightarrow C^\infty(\pi_2)$$

be $C^\infty(M)$-linear. Then φ is equivalent with a vectorbundle morphism

$$\widehat{\phi} : E_1 \longrightarrow E_2$$

over

$$\phi = Id : B \longrightarrow B.$$

From this lemma we conclude that Γ also can be considered as a morphism of vector bundles $\Gamma : \tau_M \longrightarrow \tau_M^*$.

$$\begin{array}{rccll} \Gamma : & T_xM & \longrightarrow & T_x^*M & \\ & X & \longmapsto & i_X\Omega_x & = \Omega_x(X, \bullet) \end{array}$$

over $Id : M \longrightarrow M$. One can show that this is an isomorphism.

2.2 Contact Manifolds

Let M be a $(2n+1)$-dimensional smooth manifold. A 1-form $\omega \in \Omega^1(M)$ is said to be **nondegenerate** if the top form $\omega \wedge (d\omega)^n \in \Omega^{2n+1}(M)$ is nonzero.

Definition 2.10 *Let x be a point in M and let us denote a neighborhood in M containing x by $\mathcal{O}_x$. A distribution $\mathcal{C}$ on M is said to be a **contact distribution** if there for all points $x \in M$ exists a nondegenerate 1-form $\omega \in \Omega^1(\mathcal{O}_x)$ such that $\mathcal{C} = \ker(\omega)$.*

In this case $\mathcal{C}$ is also a subbundle of τ_M and ω can be taken as a local generator of $C^\infty(\mathrm{Ann}(\mathcal{C}))$, where $\mathrm{Ann}(\mathcal{C})$ is the distribution known as the annihilator of $\mathcal{C}$. $\mathrm{Ann}(\mathcal{C})$ is a subbundle of τ_M^* with rank 1 since $\mathcal{C}$ has rank $2n$.

Definition 2.11 *A smooth manifold M together with a contact distribution $\mathcal{C} \subset \tau_M$ is said to be a **contact manifold**.*

Definition 2.12 *A smooth manifold M together with a nondegenerate 1-form $\omega \in \Omega^1(M)$ is said to be a **strict contact manifold**.*

Since every nondegenerate 1-form $\omega \in \Omega^1(M)$ defines a contact distribution, any strict contact manifold is also a contact manifold. The opposite is not true. A contact distribution $\mathcal{C}$ does not define a unique nondegenerate 1-form ω. One can easily check that if ω annihilates $\mathcal{C}$, then so does $f \cdot \omega$ for any $f \in C^\infty(M)$.

Example 2.13 *Let $M = \mathbb{R}^{2n+1}(q_1, \ldots, q_n, u, p_1, \ldots, p_n)$. Then*

$$TM \simeq \mathbb{R}^{2n+1}(\partial_{q_1}, \ldots, \partial_{q_n}, \partial_u, \partial_{p_1}, \ldots, \partial_{p_n})$$

and

$$T^*M \simeq \mathbb{R}^{2n+1}(dq_1, \ldots, dq_n, du, dp_1, \ldots, dp_n).$$

The distribution

$$\mathcal{C} = \langle \partial_{q_1} + p_1\partial_u, \ldots, \partial_{q_n} + p_n\partial_u, \partial_{p_1}, \ldots, \partial_{p_n} \rangle$$

*is a contact distribution on M. $\mathcal{C}$ is often referred to as the **Cartan distribution**. One can check that the 1-form*

$$\omega = du - p_1 dq_1 - \cdots - p_n dq_n$$

annihilates $\mathcal{C}$ and that

$$\omega \wedge (d\omega)^n = dq_1 \wedge dp_1 \wedge dq_2 \wedge dp_2 \wedge \cdots \wedge dp_n \wedge du \neq 0,$$

*hence ω is nondegenerate. This ω we call the **Cartan form**.*

Definition 2.14 *Let $\Pi \subset \tau_M$ be a distribution. A submanifold $N \subset M$ is said to be an **integral manifold** of Π if $T_xN \subset \Pi_x$ for all points x in N.*

Definition 2.15 *Let (M, Π) be a contact manifold. An integral manifold N of Π is called* ***Legendrian*** *if*

$$2 \dim N = \dim M - 1.$$

Lemma 2.16 *Let (M, ω) be a strict contact manifold and $\mathcal{C}$ the contact distribution defined by ω, i.e. $\mathcal{C} = \ker(\omega)$. Then there exists a canonical "symplectic" form Ω on $\mathcal{C}$ induced by ω.*

Proof. This follows from the fact that ω is nondegenerate. That is:

$$\omega \wedge (d\omega)^n \neq 0$$

if $\dim(M) = 2n + 1$. Let

$$\tau_M = \left\langle \xi_1, \ldots, \xi_{2n+1} \right\rangle \qquad \text{and} \qquad \mathcal{C} = \langle \xi_1, \ldots, \xi_{2n} \rangle .$$

Then the following inequality must hold:

$$\left(\omega \wedge (d\omega)^n\right) \left(\xi_1, \cdots, \xi_{2n+1}\right) \neq 0$$

Since $\mathcal{C} = \ker(\omega)$, this is equivalent to

$$\omega\left(\xi_{2n+1}\right) \cdot (d\omega)^n (\xi_1, \cdots, \xi_{2n}) \neq 0,$$

which implies that $\Omega = d\omega|_{\mathcal{C}} \neq 0$. Thus Ω_x is a canonical symplectic structure on $\mathcal{C}_x$ for all points $x \in M$. ■

2.3 Jet bundles π_k

We will now consider another important construction from the filtration of ideals

$$C^\infty(M) \supset \mu_x \supset \mu_x^2 \supset \mu_x^3 \supset \cdots$$

introduced above. From this we defined the cotangent space of the manifold M as the quotient $\frac{\mu_x}{\mu_x^2}$. Once more we are interested in quotients.

Definition 2.17 *We call the quotient*

$$J_x^k M = \frac{C^\infty(M)}{\mu_x^{k+1}}$$

the space of k-jets of functions at x in M.

Since $C^\infty(M)$ is an $\mathbb{R}$-algebra and μ_x^{k+1} is an ideal in $C^\infty(M)$, it is well known that $J_x^k M$ must be an $\mathbb{R}$-algebra. Let $\lambda \in \mathbb{R}$, $[f]_x^k, [g]_x^k \in J_x^k M$. Then one can check that

$$\lambda \cdot [f]_x^k = [\lambda f]_x^k \qquad [f]_x^k + [g]_x^k = [f+g]_x^k \qquad [f]_x^k \cdot [g]_x^k = [f \cdot g]_x^k$$

gives $J_x^k M$ an $\mathbb{R}$-algebra structure.

Definition 2.18 *We call the bundle*

$$\pi_k : J^k M \longrightarrow M$$

the k-jet bundle over M. *Here $J^k M = \bigcup_{x \in N} J_x^k M$ is a smooth manifold.*

Every function $f \in C^\infty(M)$ defines a section in π_k. We denote the map relating functions and sections by j_k.

$$\begin{array}{rccc} j_k : & C^\infty(M) & \longrightarrow & C^\infty(\pi_k) \\ & f & \longmapsto & j_k(f) = [f]^k \end{array},$$

where

$$\begin{array}{rccc} [f]^k : & M & \longrightarrow & J^k M \\ & x & \longmapsto & [f]_x^k = f \bmod \mu_x^{k+1} \end{array}.$$

We say that $j_k(f)$ is the k-jet of the function f.

Remark 2.19 *$J^1 M$ is a strict contact manifold.*

2.4 Monge-Ampere equations and effective forms

The classical Monge-Ampere equation in two variables is a second order partial differential equation with two variables q_1, q_2 and one unknown function $u(q_1, q_2)$:

$$F_1 + F_2 \frac{\partial^2 u}{\partial q_1^2} + F_3 \frac{\partial^2 u}{\partial q_2^2} + F_4 \frac{\partial^2 u}{\partial q_1 \partial q_2} + F_5 \left(\frac{\partial^2 u}{\partial q_1^2} \frac{\partial^2 u}{\partial q_2^2} - \left(\frac{\partial^2 u}{\partial q_1 \partial q_2} \right)^2 \right) = 0, \tag{2.1}$$

where $F_1, \ldots, F_5$ are functions of $q_1, q_2, u, \frac{\partial u}{\partial q_1}, \frac{\partial u}{\partial q_2}$.

From [Ly1] we have the following observation: Every differential n-form $\theta \in \Omega^n(J^1 M)$ defines a second order differential operator

$$\begin{array}{rccc} \Delta_\theta : & C^\infty(M) & \longrightarrow & \Omega^n(M) \\ & f & \longmapsto & j_1(f)^*(\theta) = \theta|_{j_1(h)} \end{array}$$

called the Monge-Ampere operator. Here $j_1(f)^*$ denotes morphism

$$j_1(f)^* : \Omega^n\left(J^1 M\right) \longrightarrow \Omega^n(M)$$

of $C^\infty(M)$-modules induced from the section $j_1(f) : M \longrightarrow J^1M$. The kernel of Δ_θ is equivalent to a second order partial differential equation in n variables.

Example 2.20 (Linear wave 1) *Let $M = \mathbb{R}^2(q_1, q_2)$. Then $J^1M \simeq \mathbb{R}^5(q_1, q_2, u, p_1, p_2)$. Consider the form*

$$\theta = dq_1 \wedge dp_2 + dq_2 \wedge dp_1.$$

The corresponding Monge-Ampere operator has the form

$$\Delta_\theta(h) = \left(-\frac{\partial^2 h}{\partial q_1^2} + \frac{\partial^2 h}{\partial q_2^2}\right) dq_1 \wedge dq_2.$$

So the Monge-Ampere equation $\Delta_\theta(h) = 0$ becomes the linear wave equation:

$$\frac{\partial^2 h}{\partial q_2^2} - \frac{\partial^2 h}{\partial q_1^2} = 0.$$

Example 2.21 (Laplace) *Similar to the preceding example one can check that the form*

$$\theta = dq_1 \wedge dp_2 + dp_1 \wedge dq_2$$

corresponds to the Laplace equation

$$\frac{\partial^2 h}{\partial q_1^2} + \frac{\partial^2 h}{\partial q_2^2} = 0.$$

Definition 2.22 *Let M be an n-dimensional smooth manifold. We call*

$$\Delta_\theta(f) = 0 \quad ; \quad \theta \in \Omega^n M$$

*the **Monge-Ampere equation**. A solution of this equation is a Legendrian submanifold $L \subset J^1M$, such that $\theta|_L = 0$ and $\pi_1|_L$ is a diffeomorphism.*

To allow multivalued solutions, the following generalization is made:

Definition 2.23 *We define **generalized solutions** of Monge-Ampere equations to be Legendrian submanifolds $L \subset J^1M$, such that $\theta|_L = 0$.*

Example 2.24 *Let $M = \mathbb{R}^2(q_1, q_2)$. Then one can show that the form*

$$\theta = F_1 dq_1 \wedge dq_2 + F_2 dq_2 \wedge dp_1 + F_3 dq_1 \wedge dp_2 + F_4 dq_1 \wedge dp_1 - F_4 dq_2 \wedge dp_2 + F_5 dp_1 \wedge dp_2,$$

where the coefficients are functions in q_1, q_2, u, p_1 and p_2, corresponds to the Monge-Ampere equation

$$F_1 - F_2 \frac{\partial^2 u}{\partial q_1^2} + F_3 \frac{\partial^2 u}{\partial q_2^2} + 2F_4 \frac{\partial^2 u}{\partial q_1 \partial q_2} + F_5 \left(\frac{\partial^2 u}{\partial q_1^2}\frac{\partial^2 u}{\partial q_2^2} - \left(\frac{\partial^2 u}{\partial q_1 \partial q_2}\right)^2\right) = 0.$$

Although a given form θ defines an unique Monge-Ampere operator Δ_θ, the correspondence is not one-to-one. The correspondence is presented in [Ly1] and [Ly2]. We present the main steps of argumentation for the sake of completeness. We investigate the graded, super commutative, differential $\mathbb{R}$-algebra

$$\Omega^* \left(J^1 M\right) = \bigoplus_{i=0}^{\dim J^1 M} \Omega^i \left(J^1 M\right).$$

Let ω denote the contact form in correspondence with the strict contact manifold $J^1 M$, i.e. $\omega \in \Omega^1 \left(J^1 M\right)$ is nondegenerate.

Theorem 2.25 (Lepage) *Two k'th order Monge-Ampere operators Δ_θ and $\Delta_{\theta'}$ are equal if and only if the differential k-form $\theta' \in \Omega^k \left(J^1 M\right)$ can be written in the form:*

$$\theta' = \theta + \alpha \wedge \omega + \beta \wedge d\omega$$

for some differential forms $\alpha \in \Omega^{k-1} \left(J^1 M\right)$, $\beta \in \Omega^{k-2} \left(J^1 M\right)$.

Proof. This statement with proof can be found in [Ly1] and [Ly2]. ■

This theorem leads to the following definition.

Definition 2.26 *Let I^k be a subspace of $\Omega^k \left(J^1 M\right)$ defined by:*

$$I^k = \left\{ \mu \in \Omega^k \left(J^1 M\right) \middle| \, \mu = \alpha \wedge \omega + \beta \wedge d\omega \text{ for some } \alpha \in \Omega^{k-1} \left(J^1 M\right), \beta \in \Omega^{k-2} \left(J^1 M\right) \right\}$$

Then we call the ideal

$$I_{\mathcal{C}} = \bigoplus_k I^k$$

*in the exterior algebra $\Omega^* \left(J^1 M\right)$, the **Cartan ideal**.*

It is easy to see that if $\mu \in I^k$, then $d\mu \in I^{k+1}$, so $I_{\mathcal{C}}$ is a graded, super commutative, differential ideal in $\Omega^* \left(J^1 M\right)$. Thus the factor $\Omega^* \left(J^1 M\right) \diagup I_{\mathcal{C}}$ must be a graded, super commutative, differential $\mathbb{R}$-algebra.

Definition 2.27 *We define **the algebra of effective forms** to be*

$$\Omega^*_{\text{eff}} \left(J^1 M\right) = \frac{\Omega^* \left(J^1 M\right)}{I_{\mathcal{C}}}.$$

The Lepage theorem gives us a one-to-one correspondence between effective forms and Monge-Ampere operators Δ_θ. But it still remains to define Monge-Ampere operators from effective forms. We need to find a canonical representative θ_0 in $\Omega^* \left(J^1 M\right)$ from any given effective form $\theta_{\text{eff}} \in \Omega^*_{\text{eff}} \left(J^1 M\right)$. Then we can define $\Delta_{\theta_{\text{eff}}}$ to be Δ_{θ_0}.

We know that the contact form ω induces a contact distribution $\mathcal{C} \subset \tau_{J^1M}$ and the symplectic structure $\Omega = d\omega|_{\mathcal{C}}$. Ω implies the isomorphism

$$\Gamma : \mathcal{C} \longrightarrow \mathcal{C}^*$$

as mentioned above. The exact sequences

$$0 \longrightarrow \mathcal{C} \longrightarrow \tau_{J^1M} \longrightarrow \left(\frac{\tau_{J^1M}}{\mathcal{C}}\right) \longrightarrow 0$$

and

$$0 \longrightarrow \mathcal{C}^* \longrightarrow \tau^*_{J^1M} \longrightarrow \left(\frac{\tau_{J^1M}}{\mathcal{C}}\right)^* \longrightarrow 0$$

are well known. Using that $\left(\frac{\tau_{J^1M}}{\mathcal{C}}\right) \simeq \mathrm{Ann}\,(\mathcal{C}^*)$ and $\left(\frac{\tau_{J^1M}}{\mathcal{C}}\right)^* \simeq \mathrm{Ann}\,(\mathcal{C})$ we get the isomorphisms

$$\tau^*_{J^1M} \simeq \mathcal{C}^* \oplus \mathrm{Ann}\,(\mathcal{C}) \qquad \text{and} \qquad \tau_{J^1M} \simeq \mathrm{Ann}\,(\mathcal{C}^*) \oplus \mathcal{C}$$

These isomorphisms imply

$$\begin{aligned} \Lambda^k \tau^*_{J^1M} &\simeq \Lambda^k \mathcal{C}^* \oplus \Lambda^{k-1}\mathcal{C}^* \wedge \mathrm{Ann}\,(\mathcal{C}), \\ \Lambda^k \tau_{J^1M} &\simeq \mathrm{Ann}\,(\mathcal{C}^*) \wedge \Lambda^{k-1}\mathcal{C} \oplus \Lambda^k \mathcal{C}. \end{aligned}$$

Since the contact form ω can be taken as the generator of $\mathrm{Ann}\,(\mathcal{C})$, this leads to the isomorphism

$$\Omega^k_{\mathrm{eff}}\left(J^1M\right) \simeq \frac{C^\infty\left(\Lambda^k \mathcal{C}^*\right)}{\left\{\beta \wedge d\omega|_{\mathcal{C}} \,|\, \beta \in C^\infty\left(\Lambda^{k-2}\mathcal{C}^*\right)\right\}}.$$

Let a be a point in J^1M. Let us investigate the space $\Lambda^k\left(C^*_a\right)$. We introduce the two operators

$$\begin{array}{rccc} \top: & \Lambda^i \mathcal{C}^*_a & \longrightarrow & \Lambda^{i+2}\mathcal{C}^*_a \\ & \theta & \longmapsto & \theta \wedge \Omega \end{array}$$

and

$$\begin{array}{rccc} \bot: & \Lambda^i \mathcal{C}^*_a & \longrightarrow & \Lambda^{i-2}\mathcal{C}^*_a \\ & \theta & \longmapsto & i_{X_\Omega}\theta \end{array},$$

where X_Ω denotes the canonical bivector field $\Lambda^2\Gamma^{-1}(\Omega)$.

Theorem 2.28 (Hodge-Lepage) *For any k-form $\theta \in \Lambda^k \mathcal{C}^*_a$, the following Hodge-Lepage decomposition holds:*

$$\theta = \theta_0 + \top\theta_1 + \top^2\theta_2 + \cdots,$$

where $\theta_i \in \Lambda^{k-2i}\mathcal{C}^*_a$ are uniquely determined forms satisfying $\bot\theta_i = 0$.

Proof. This statement with proof can be found in [Ly1] and [Ly2]. ■

From [Ly1] and [Ly2] we have an explicit expression for θ_0:

$$\theta_0 = \left(\sum_{i \geq 0} (-1)^i \frac{1}{(i+1)!i!} \top^i \perp^i \right) (\theta) .$$

So we have the isomorphism

$$\begin{aligned} \Omega^k_{\text{eff}} \left(J^1 M \right) &\simeq \left\{ \mu \in C^\infty \left(\Lambda^k \mathcal{C}^* \right) \middle| \perp \mu = 0 \right\} \stackrel{\text{def}}{=} C^\infty \left(\Lambda^k_{\text{eff}} \mathcal{C}^* \right), \\ \theta \bmod I^k &\simeq \theta_0 = \left(\sum_{i \geq 0} (-1)^i \frac{1}{(i+1)!i!} \top^i \perp^i \right) (\theta) . \end{aligned}$$

This give us the one-to-one correspondence between Monge-Ampere operators of k'th order and effective forms $\theta_{\text{eff}} \in C^\infty \left(\Lambda^k_{\text{eff}} C^* \right)$.

Lemma 2.29 *A Monge-Ampere equation in k variables, defines a conformal effective form $\theta \in C^\infty \left(\Lambda^k_{\text{eff}} \mathcal{C}^* \right)$.*

Proof. We know the equivalence between effective forms $\theta_{\text{eff}} \in C^\infty \left(\Lambda^k_{\text{eff}} \mathcal{C}^* \right)$ and Monge-Ampere operators $\Delta_{\theta_0} : C^\infty (M) \longrightarrow \Omega^k (M)$. Remember that a Monge-Ampere equation is equivalent to the kernel of a Monge-Ampere operator:

$$\Delta_{\theta_0} (f) = 0.$$

Let $g \in \Omega^0 (J^1 M) = C^\infty (J^1 M)$. Then we know that

$$\Delta_{g \cdot \theta_0} (f) = j_1 (f)^* (g \cdot \theta_0) = j_1 (f)^* (g) \cdot j_1 (f)^* (\theta_0) = (g \circ j_1 (f)) \cdot \Delta_{\theta_0} (f) .$$

So we see that if g is a nonzero function, then

$$\ker (\Delta_{g \cdot \theta_0}) = \ker (\Delta_{\theta_0}) .$$

The claim follows. ■

Let $\mathbb{R}^* = \mathbb{R} \backslash \{0\}$. Then the Monge-Ampere equation $\Delta_\theta (f) = 0$ defines at every point a in $J^1 M$ the subset $\mathbb{R}^* \cdot \theta_0(a)$ in $\Lambda^k \mathcal{C}^*_a$, where θ_0 is the effective part of θ according to the Hodge-Lepage decompositon.

2.5 Classification of effective forms

Our main object in this thesis will be a 5-dimensional strict contact manifold. This corresponds to $J^1 M$, where M is 2-dimensional. Hence this is related to second order PDE's of Monge-Ampere type with 2 variables. A PDE like this corresponds to a conformal effective form $\theta \in C^\infty (\Lambda^2_{\text{eff}} \mathcal{C}^*)$. The following proposition from [Ly1] gives us a useful description of the space $C^\infty (\Lambda^2_{\text{eff}} \mathcal{C}^*)$.

Proposition 2.30 *When the Cartan distribution $\mathcal{C}$ has rank $2n$, the form $\omega \in \Lambda^{n-k}\mathcal{C}_a^*$ is effective if and only if $\top^{k+1}(\omega) = 0$.*

When $\mathcal{C}$ has rank 4, it follows that $\{\mu \in \Lambda^2\mathcal{C}^* | \top\mu = 0\} = \{\mu \in \Lambda^2\mathcal{C}^* | \perp\mu = 0\}$, so we can write

$$C^\infty\left(\Lambda^2_{\text{eff}}\mathcal{C}^*\right) = \left\{\mu \in C^\infty\left(\Lambda^2\mathcal{C}^*\right) \middle| \mu \wedge \Omega = 0\right\},$$

where $\Omega \in C^\infty(\Lambda^2\mathcal{C}^*)$ is the symplectic structure in $\mathcal{C}$.

In [Ly2] there is given a pointwise classification of effective forms $\theta \in C^\infty(\Lambda^2_{\text{eff}}\mathcal{C}^*)$. Let a be a point in J^1M and let $\Omega_a \in \Lambda^2\mathcal{C}_a^*$ be the symplectic form. Then by dimensional reasons $\theta_a \wedge \theta_a$ and $\Omega_a \wedge \Omega_a$ are proportional.

Definition 2.31 *We define the **Pfaffian** of a 2-form θ_a in $\Lambda^2\mathcal{C}_a^*$ by:*

$$\theta_a \wedge \theta_a = \text{Pf}(\theta_a)\,\Omega_a \wedge \Omega_a,$$

where Ω_a is the symplectic form in $\mathcal{C}_a$.

Definition 2.32 *An effective 2-form $\theta_a \in \Lambda^k\mathcal{C}_a^*$ is said to be*

- ***elliptic** if $\text{Pf}(\theta_a) > 0$,*
- ***hyperbolic** if $\text{Pf}(\theta_a) < 0$,*
- ***parabolic** if $\text{Pf}(\theta_a) = 0$ and $\theta_a \neq 0$.*

Example 2.33 *Let $M = \mathbb{R}^2(q_1, q_2)$ and $J^1M \simeq \mathbb{R}^5(q_1, q_2, u, p_1, p_2)$. Then one can show ([Ly2]) that an effective 2-form θ must have the form:*

$$\theta = F_1 dq_1 \wedge dq_2 + F_2 dq_2 \wedge dp_1 + F_3 dq_1 \wedge dp_2 + F_4 dq_1 \wedge dp_1 - F_4 dq_2 \wedge dp_2 + F_5 dp_1 \wedge dp_2.$$

And when $\Omega = dq_1 \wedge dp_1 + dq_2 \wedge dp_2$ is the symplectic form, then

- *θ is hyperbolic when $F_1F_5 + F_2F_3 + F_4^2 > 0$,*
- *θ is elliptic when $F_1F_5 + F_2F_3 + F_4^2 < 0$,*
- *θ is parabolic when $F_1F_5 + F_2F_3 + F_4^2 = 0$.*

We will consider the pure hyperbolic case, that is when $\text{Pf}(\theta_a) < 0 \;\forall\; a \in J^1M$. For short we will write $\text{Pf}(\theta) < 0$.

With any 2-form $\theta_a \in \Lambda^k\mathcal{C}_a^*$ we may associate an operator

$$j : \mathcal{C}_a \longrightarrow \mathcal{C}_a$$

by requiring that $\theta_a(X, Y) = \Omega_a(jX, Y)$ holds for all vectors X, Y in $\mathcal{C}_a$ ([LRC]). From the bilinearity of θ_a and Ω_a it follows that j is a linear operator.

Definition 2.34 *Let (M, ω) be a 5-dimensional strict contact manifold. Let Π be the contact distribution induced by ω, and let $\theta \in C^\infty(\Lambda^2 \Pi^*)$ be effective. Then we say that (ω, θ) is a* ***generalized Monge-Ampere equation.*** *(ω, θ) is elliptic, hyperbolic or parabolic according to the effective form θ.*

Proposition 2.35 *For any effective 2-form $\theta_a \in \Lambda^k \mathcal{C}_a^*$ the equality*

$$j^2 = -\operatorname{Pf}(\theta_a)$$

holds.

Proof. The statement and proof can be found in [Ly2]. ■

In the hyperbolic case we see that $j^2 = |\operatorname{Pf}(\theta_a)| \neq 0$, so j is invertible, $j^{-1} = \frac{1}{|\operatorname{Pf}(\theta_a)|} j$, and hence it is a linear transformation of $\mathcal{C}_a$.

Example 2.36 (General j) *Let $M = \mathbb{R}^2(q_1, q_2)$ and $J^1 M \simeq \mathbb{R}^5(q_1, q_2, u, p_1, p_2)$. In the case*

$$\theta = F_1 dq_1 \wedge dq_2 + F_2 dq_2 \wedge dp_1 + F_3 dq_1 \wedge dp_2 + F_4 dq_1 \wedge dp_1 - F_4 dq_2 \wedge dp_2 + F_5 dp_1 \wedge dp_2,$$
$$\Omega = dq_1 \wedge dp_1 + dq_2 \wedge dp_2,$$
$$\mathcal{C} = \langle \partial_{q_1} + p_1 \partial_u, \partial_{q_2} + p_2 \partial_u, \partial_{p_1}, \partial_{p_2} \rangle = \langle \mathcal{D}_{q_1}, \mathcal{D}_{q_2}, \partial_{p_1}, \partial_{p_2} \rangle,$$

then j has the form

$$j = \begin{bmatrix} F_4 & F_2 & 0 & -F_5 \\ F_3 & -F_4 & F_5 & 0 \\ 0 & F_1 & F_4 & F_3 \\ -F_1 & 0 & F_2 & -F_4 \end{bmatrix}$$

in the basis $\{\mathcal{D}_{q_1}, \mathcal{D}_{q_2}, \partial_{p_1}, \partial_{p_2}\}$ into $\mathcal{C}$.

Notation 2.37 *In later examples we will consequently write $\mathcal{D}_{q_1}$ and $\mathcal{D}_{q_2}$ instead of $\partial_{q_1} + p_1 \partial_u$ and $\partial_{q_2} + p_2 \partial_u$.*

Chapter 3

Hyperbolic M-A equations and 5-dimensional strict contact manifolds

In this chapter we discover two canonical rank 1 distributions, L^1 and χ, in τ_M. These subbundles imply two canonical decompositions of τ_M. In section 3 we introduce and investigate the Nijenhuis tensor N_j.

3.1 The extending sequence of derived submodules

From any distribution $\Sigma \subset \tau_M$ we can construct the so called extending sequence of derived submodules. We know that $\mathcal{D}_1(M)$ is a $C^\infty(M)$-module. Consider a neighborhood (germ) $\mathcal{O}_x$ of x in M. We say that $\xi_1, \ldots, \xi_t \in \mathcal{D}_1(M)$ are local generators of $C^\infty(\Sigma)$ (in $\mathcal{O}_x$) if $\xi_1(y), \ldots, \xi_t(y)$ is a generating set of vectors in Σ_y for any $y \in \mathcal{O}_x$ and we write

$$C^\infty(\Sigma) = \langle \xi_1, \ldots, \xi_t \rangle .$$

Define

$$\mathcal{D}_1(\Sigma) = \{ \eta \in \mathcal{D}_1(M) | \, \eta = f_1\xi_1 + \cdots + f_t\xi_t, \forall y \in \mathcal{O}_x \},$$

where $f_1, \ldots, f_t \in C^\infty(M)$. Then $\mathcal{D}_1(\Sigma)$ is a submodule of $\mathcal{D}_1(M)$. Now we define the $C^\infty(M)$-module $\mathcal{D}_{k+1}(\Sigma)$ to be generated by $\mathcal{D}_k(\Sigma)$ and all possible commutators $[\chi, \xi]$ where $(\chi, \xi) \in \mathcal{D}_k(\Sigma) \times \mathcal{D}_1(\Sigma)$.

$$\mathcal{D}_{k+1}(\Sigma) = [\mathcal{D}_k(\Sigma), \mathcal{D}_1(\Sigma)] .$$

Then we get **the extending sequence of derived submodules**:

$$\mathcal{D}_1(\Sigma) \subset \mathcal{D}_2(\Sigma) \subset \cdots \subset \mathcal{D}_1(M) .$$

In general $\mathcal{D}_k(\Sigma)$ is not a module of sections for some distribution, since the rank of its localization can vary from point to point. The module $\mathcal{D}_k(\Sigma)$ is **projective** if it is a direct summand in a free module. Then, in our situation, it is a module of sections for some distribution Σ_k. We call Σ_k the $(k-1)$-extension of Σ and we denote it $\partial^{(k-1)}\Sigma$.

How do we localize the module $\mathcal{D}_k(\Sigma)$? We define

$$\Sigma_{k,x} = \frac{\mathcal{D}_k(\Sigma)}{\mu_x \mathcal{D}_k(\Sigma)},$$

where μ_x is the ideal $\{ f \in C^\infty(M) | f(x) = 0\}$ as before. It is well known that

$$T_x M \simeq \frac{\mathcal{D}_1(M)}{\mu_x \mathcal{D}_1(M)},$$

so we see that $\Sigma_{k,x}$ is a subspace of $T_x M$. Thus we get a sequence of vectorbundles:

$$\Sigma_1 \subseteq \Sigma_2 \subseteq \cdots \subseteq \tau_M.$$

From these vectorbundles we can define the associated vectorbundles

$$\Sigma^{[k+1]} = \frac{\Sigma_{k+1}}{\Sigma_k}$$

and get the graded bundle

$$\Sigma_1 \oplus \Sigma^{[2]} \oplus \cdots \oplus \Sigma^{[k]}$$

associated to the flag of $C^\infty(M)$-modules. This graded bundle is locally isomorphic to the tangent bundle τ_M when $\mathcal{D}_k(\Sigma) = \mathcal{D}_1(M)$ for some k.

Example 3.1 *Consider $M = \mathbb{R}^3(x,u,p)$. Let $\Pi = \ker(du - pdx) = \langle \partial_x + p\partial_u, \partial_p \rangle$. Since $[\partial_p, \partial_x + p\partial_u] = \partial_u \notin \Pi$, we have $\mathcal{D}_2 = \mathcal{D}_1(M)$. Thus we can identify $\Pi^{[2]} = \frac{\Pi_2}{\Pi} = \langle \partial_u \rangle$ and $TM \simeq \Pi \oplus \Pi^{[2]} = \langle \partial_x + p\partial_u, \partial_p \rangle \oplus \langle \partial_u \rangle$.*

3.2 Two canonical decompositions of τ_M

Let (M, ω) be a 5-dimensional strict contact manifold. We know that ω defines a rank 4 contact distribution Π and a symplectic structure $\Omega = d\omega|_\Pi$ upon Π. Let $\langle \theta \rangle$ be a conformal, hyperbolic, effective 2-form in $\Lambda^2 \Pi^*$. We get a unique representative $\theta \in \langle \theta \rangle$ by requiring $\mathrm{Pf}(\theta) = -1$. From θ and Ω we define the automorphism

$$j : \Pi \longrightarrow \Pi$$

by requiring $\theta_x(X,Y) = \Omega_x(jX, Y)$ $\forall X, Y \in \Pi_x$ $\forall x \in M$. From proposition 2.35 it follows that

$$j^2 = -\mathrm{Pf}(\theta) = \mathbf{1},$$

so j is an almost product structure on Π. From the nondegenracy and effectivity of θ one can easily show that j must have the spectrum

$$\operatorname{spec}(j) = \{1, 1, -1, -1\},$$

and it follows from the nondegeneracy of θ and Ω that

$$\dim E_j(1) = \dim E_j(-1) = 2,$$

where E_j denotes the eigenspaces of j. Hence the vector space Π_x splits into $E_j(1) \oplus E_j(-1)$ at the point x. This leads to the lemma.

Lemma 3.2 *The automorphism $j : \Pi \longrightarrow \Pi$ splits the rank 4 distribution Π into the direct sum of two rank 2 distributions:*

$$\Pi = \Pi^2_+ \oplus \Pi^2_-,$$

where

$$\Pi^2_+ = \{X \in \Pi \mid jX = X\} = \ker(j - 1)$$

and

$$\Pi^2_- = \{X \in \Pi \mid jX = -X\} = \ker(j + 1).$$

So any vector $X \in \Pi_x$ can be written as $X_+ + X_- \in \Pi^2_+ \oplus \Pi^2_-$. It's easy to see that $\left(\frac{1}{2} + \frac{1}{2}j\right)$ and $\left(\frac{1}{2} - \frac{1}{2}j\right)$ are projections in Π onto Π^2_+ and Π^2_- respectively. Let us denote

$$P_+ = \frac{1}{2}(1 + j) \qquad \text{and} \qquad P_- = \frac{1}{2}(1 - j).$$

Lemma 3.3 *The distributions Π^2_+ and Π^2_- are skew orthogonal, $\Pi^2_+ \perp_\Omega \Pi^2_-$, that is*

$$\Omega(\xi, \eta) = 0 \ \forall\ \xi \in \Pi^2_+, \eta \in \Pi^2_-.$$

Proof. Let $\xi \in C^\infty\left(\Pi^2_+\right)$ and $\eta \in C^\infty\left(\Pi^2_-\right)$. So $j\xi = \xi$ while $j\eta = -\eta$. We observe that

$$\theta(\xi, \eta) = \Omega(j\xi, \eta) = \Omega(\xi, \eta) = -\Omega(\eta, \xi) = \Omega(j\eta, \xi) = \theta(\eta, \xi) = -\theta(\xi, \eta).$$

We conclude that $\Omega(\xi, \eta) = 0 \ \forall\ \xi \in \Pi^2_+,\ \eta \in \Pi^2_-$. ■

Recall the notion of integrable distributions.

Definition 3.4 *A distribution Π on the n-dimensional manifold M is said to be **integrable** if there exist coordinates $q_1, \ldots, q_n$ on M s.t. $\Pi = \langle \partial_{q_1}, \ldots, \partial_{q_r} \rangle$, $r = \operatorname{rank}(\Pi) \leq n$.*

Theorem 3.5 (Frobenius) *A distribution $\Pi \subset \tau_M$, generated by $\xi_1, \ldots, \xi_k \in \mathcal{D}_1(M)$ is integrable if*

$$\left[\xi_i, \xi_j\right] \in C^\infty(\Pi) \quad \forall i, j.$$

Proof. The theorem and proof is presented in [S]. ■

Lemma 3.6 *The distributions Π, Π_+^2 and Π_-^2 are nonintegrable.*

Proof. This proof follows from the Cartan formula for 2-forms:

$$d\omega(\xi, \eta) = \xi(\omega(\eta)) - \eta(\omega(\xi)) - \omega([\xi, \eta]). \tag{3.1}$$

Here $\omega \in \Omega^1(M)$ is our contact form and $\xi, \eta \in \mathcal{D}_1(M)$. Let $\xi, \eta \in \mathcal{D}_1(M)$ be tangent to Π. Then the Cartan formula restricted to Π becomes

$$\Omega(\xi, \eta) = -\omega([\xi, \eta]). \tag{3.2}$$

Assume that $\xi \in \mathcal{D}_1(M)$ is tangent to Π_+^2. Ω is nondegenerate on Π, so there exist some $\eta \in \mathcal{D}_1(M)$ that is tangent to Π such that $\Omega(\xi, \eta) \neq 0$. Since $\Omega\left(\Pi_+^2, \Pi_-^2\right) = 0$, η cannot be tangent to Π_-^2. Thus if we decompose η according to the direct sum $C^\infty(\Pi) \simeq C^\infty\left(\Pi_+^2\right) \oplus_{C^\infty(M)} C^\infty\left(\Pi_-^2\right)$, that is

$$\eta = \eta_+ + \eta_- \in C^\infty\left(\Pi_+^2\right) \oplus_{C^\infty(M)} C^\infty\left(\Pi_-^2\right),$$

then η_+ must be nonzero. Hence

$$\Omega(\xi, \eta) = \Omega(\xi, \eta_+) = -\omega(\left[\xi, \eta_+\right]) \neq 0.$$

This implies that $\left[\xi, \eta_+\right]$ cannot be tangent to Π. We conclude that Π and Π_+^2 are nonintegrable. Similarly we can prove that Π_-^2 is nonintegrable. ■

Corollary 3.7 *$\Omega|_{\Pi_+^2}$ and $\Omega|_{\Pi_-^2}$ are nondegenerate, hence $\Pi_\pm^2 \subset \Pi$ are symplectic subbundles.*

Corollary 3.8 *$\left[\xi_\pm, \eta_\mp\right] \in C^\infty(\Pi)$ for all sections $\xi_\pm, \eta_\pm \in C^\infty\left(\Pi_\pm^2\right)$.*

Corollary 3.9 *There exists sections $\xi_\pm, \eta_\pm \in C^\infty\left(\Pi_\pm^2\right)$ such that $\left[\xi_\pm, \eta_\pm\right] \notin C^\infty(\Pi)$.*

The modules

$$\left[C^\infty\left(\Pi_+^2\right), C^\infty\left(\Pi_+^2\right)\right] \quad \text{and} \quad \left[C^\infty\left(\Pi_-^2\right), C^\infty\left(\Pi_-^2\right)\right]$$

are projective since $\left(\Pi_+^2, \Omega|_{\Pi_+^2}\right)$ and $\left(\Pi_-^2, \Omega|_{\Pi_-^2}\right)$ are 2-dimensional and "symplectic". Hence there exist distributions $\partial\Pi_+^2, \partial\Pi_-^2 \subset \tau_M$ such that

$$C^\infty\left(\partial\Pi_\pm^2\right) = \left[C^\infty\left(\Pi_\pm^2\right), C^\infty\left(\Pi_\pm^2\right)\right].$$

The extensions $\partial\Pi^2_\pm$ must be of rank 3. Let us denote them

$$\Pi^3_\pm = \partial\Pi^2_\pm. \tag{3.3}$$

From corollary 3.9 it follows that $\Pi^3_\pm \not\subseteq \Pi$.

Proposition 3.10 *The distributions $\Pi^2_\pm$ and $\Pi^3_\mp$ have zero intersection:*

$$\Pi^2_\pm \cap \Pi^3_\mp = 0. \tag{3.4}$$

Proof. Let ξ_1, ξ_2 and η_1, η_2 in $\mathcal{D}_1(M)$ be generators of $C^\infty\left(\Pi^2_+\right)$ and $C^\infty\left(\Pi^2_-\right)$ in the neighborhood $\mathcal{O} \subset M$ containing x. Due to nonintegrability of Π^2_+ we know that $[\xi_1, \xi_2] \notin C^\infty(\Pi)$ in $\mathcal{O}$ for ξ_1, ξ_2 to be well defined as generators. Then $\xi_1, \xi_2, \xi_3 = [\xi_1, \xi_2]$ are local generators of $C^\infty\left(\partial\Pi^2_+\right)$. From this we get the following bases into the fibers $\Pi^2_{+,x}, \partial\Pi^2_{+,x}$ and $\Pi^2_{-,x}$:

$$\begin{aligned}
\Pi^2_{+,x} &= \operatorname{span}\{\xi_1(x), \xi_2(x)\} = \operatorname{span}\{X_1, X_2\}, \\
\partial\Pi^2_{+,x} &= \operatorname{span}\{\xi_1(x), \xi_2(x), \xi_3(x)\} = \operatorname{span}\{X_1, X_2, X_3\}, \quad \text{where } X_3 \notin \Pi_x, \\
\Pi^2_{-,x} &= \operatorname{span}\{\eta_1(x), \eta_2(x)\} = \operatorname{span}\{Y_1, Y_2\}.
\end{aligned}$$

Assume that $\partial\Pi^2_{+,x}$ and $\Pi^2_{-,x}$ have nonzero intersection, that is: $aX_1 + bX_2 + cX_3 \in \Pi^2_{-,x}$ for some $a, b, c \in \mathbb{R}$. For $aX_1 + bX_2 + cX_3$ to lie in Π_x we must require that $c = 0$ since Π_x is closed under addition. Hence $aX_1 + bX_2 + cX_3 \in \Pi^2_{-,x} \subset \Pi_x$ implies that $aX_1 + bX_2 = dY_1 + eY_2$. This contradicts the fact that $\Pi^2_{+,x} \cap \Pi^2_{-,x} = 0$, thus our assumption must be wrong. We conclude that $\partial\Pi^2_{+,x} \cap \Pi^2_{-,x} = 0$ for all x in $\mathcal{O}$. Since $\mathcal{O}$ is not specified we conclude that $\Pi^3_+ \cap \Pi^2_- = 0$. Similarly we can prove that $\Pi^3_- \cap \Pi^2_+ = 0$. ■

Corollary 3.11 $\Pi^3_+(x) \cap \Pi^3_-(x) \not\subseteq T_xM$ *and* $\dim(\Pi^3_+(x) \cap \Pi^3_-(x)) = 1$.

Proof. Obvious since T_xM is 5-dimensional and $\Pi^2_+ \cap \Pi^2_- = 0$. ■

Definition 3.12 *We define the rank 1 vectorbundle* $L^1 = \bigcup_{x \in M} \Pi^3_+(x) \cap \Pi^3_-(x)$.

So we have a canonical decomposition of τ_M:

$$\tau_M = \Pi \oplus L^1 = \Pi^2_+ \oplus \Pi^2_- \oplus L^1. \tag{3.5}$$

Example 3.13 (Linear wave 2) *Consider $M = \mathbb{R}^5(q_1, q_2, u, p_1, p_2)$ with the contact form $\omega = du - p_1 dq_1 - p_2 dq_2$. Then the contact distribution and symplectic form become*

$$\begin{aligned}
\Pi &= \langle \mathcal{D}_{q_1}, \mathcal{D}_{q_2}, \partial_{p_1}, \partial_{p_2} \rangle, \\
\Omega &= d\omega|_\Pi = dq_1 \wedge dp_1 + dq_2 \wedge dp_2.
\end{aligned}$$

We know that the linear wave equation

$$\frac{\partial^2 h}{\partial q_2^2} - \frac{\partial^2 h}{\partial q_1^2} = 0$$

corresponds to the hyperbolic effective form $\theta = dq_1 \wedge dp_2 + dq_2 \wedge dp_1$. *For the given basis into* Π *the operator* $j : \Pi \longrightarrow \Pi$ *has the matrix*

$$j = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix}.$$

We remember that $P_+ = \left(\frac{1}{2}j + \frac{1}{2}\right)$ *and* $P_- = \left(-\frac{1}{2}j + \frac{1}{2}\right)$ *are projections onto* Π_+^2 *and* Π_-^2 *correspondingly. These projections lead to the following bases in* Π_+^2 *and* Π_-^2*:*

$$\begin{aligned} \Pi_+^2 &= \ker P_- = \langle \mathcal{D}_{q_1} + \mathcal{D}_{q_2}, \partial_{p_1} + \partial_{p_2} \rangle, \\ \Pi_-^2 &= \ker P_+ = \langle \mathcal{D}_{q_1} - \mathcal{D}_{q_2}, \partial_{p_1} - \partial_{p_2} \rangle. \end{aligned}$$

From this we calculate the extensions:

$$\begin{aligned} \partial\Pi_-^2 &= \Pi_-^3 = \langle \mathcal{D}_{q_1} - \mathcal{D}_{q_2}, \partial_{p_1} - \partial_{p_2}, \partial_u \rangle, \\ \partial\Pi_+^2 &= \Pi_+^3 = \langle \mathcal{D}_{q_1} + \mathcal{D}_{q_2}, \partial_{p_1} + \partial_{p_2}, \partial_u \rangle. \end{aligned}$$

We see that

$$L^1 = \Pi_+^3 \cap \Pi_-^3 = \langle \partial_u \rangle.$$

Example 3.14 (Nonlinear 1) *Consider the same contact manifold as described in the preceding example. Let* $S, T \in C^\infty(M)$. *The M-A equation*

$$-S\frac{\partial^2 u}{\partial q_1^2} + 2\frac{\partial^2 u}{\partial q_1 \partial q_2} - T\left(\frac{\partial^2 u}{\partial q_1^2}\frac{\partial^2 u}{\partial q_2^2} - \left(\frac{\partial^2 u}{\partial q_1 \partial q_2}\right)^2\right) = 0$$

corresponds to the hyperbolic effective form

$$\theta = dq_1 \wedge dp_1 - dq_2 \wedge dp_2 + S dq_2 \wedge dp_1 - T dp_1 \wedge dp_2.$$

The operator j *has the matrix*

$$j = \begin{bmatrix} 1 & S & 0 & T \\ 0 & -1 & -T & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & S & -1 \end{bmatrix}.$$

By the projections P_+ and P_- we find the distributions

$$\begin{aligned}\Pi_+^2 &= \langle \mathcal{D}_{q_1}, T\mathcal{D}_{q_2} - 2\partial_{p_1} - S\partial_{p_2} \rangle, \\ \Pi_-^2 &= \langle S\mathcal{D}_{q_1} - 2\mathcal{D}_{q_2}, T\mathcal{D}_{q_2} - S\partial_{p_2} \rangle.\end{aligned}$$

The extensions become:

$$\Pi_+^3 = \langle \mathcal{D}_{q_1}, T\mathcal{D}_{q_2} - 2\partial_{p_1} - S\partial_{p_2}, \mathcal{D}_{q_1}(T)\mathcal{D}_{q_2} + 2\partial_u - \mathcal{D}_{q_1}(S)\partial_{p_2} \rangle$$

and

$$\begin{aligned}\Pi_-^3 &= \langle S\mathcal{D}_{q_1} - 2\mathcal{D}_{q_2}, T\mathcal{D}_{q_2} - S\partial_{p_2}, \left(S\frac{\partial S}{\partial p_2} - T\mathcal{D}_{q_2}(S)\right)\mathcal{D}_{q_1} + (S\mathcal{D}_{q_1}(T) - 2\mathcal{D}_{q_2}(T))\mathcal{D}_{q_2} \\ &\quad -2S\partial_u + (2\mathcal{D}_{q_2}(S) - S\mathcal{D}_{q_1}(S))\partial_{p_2} \rangle.\end{aligned}$$

One can calculate the intersection to be

$$L^1 = \left\langle \left(S\mathcal{D}_{q_1}(T) - T\mathcal{D}_{q_1}(S) - \mathcal{D}_{q_2}(T) + \frac{\partial S}{\partial p_2}\right)\mathcal{D}_{q_1} - \mathcal{D}_{q_1}(T)\mathcal{D}_{q_2} - 2\partial_u + \mathcal{D}_{q_1}(S)\partial_{p_2} \right\rangle.$$

It is well known that the contact form $\omega \in \Omega^1(M)$ defines a unique vector field in $\mathcal{D}_1(M)$ by the intersection

$$\ker(d\omega) \cap \{\xi \in \mathcal{D}_1(M) | \, i_\xi(\omega) = 1\}.$$

This vector field is said to be the **characteristic vector field** or the **Reeb vector field** of the contact structure. Let us denote it by X_1. Hence

$$i_{X_1} d\omega = 0 \qquad \text{and} \qquad \omega(X_1) = 1.$$

Obviously this vector field is not tangent to the contact distribution. X_1 generates a subbundle χ of τ_M:

$$\chi = \langle X_1 \rangle.$$

This gives us the second canonical decomposition of τ_M:

$$\tau_M = \Pi_+^2 \oplus \Pi_-^2 \oplus \chi.$$

Example 3.15 *In the case $M = \mathbb{R}^5(q_1, q_2, u, p_1, p_2)$ with the contact form $\omega = du - p_1 dq_1 - p_2 dq_2$, one can check that the characteristic vector field X_1 is $\partial_u \in \mathcal{D}_1(M)$.*

3.3 The Nijenhuis tensor

The decompositions $\tau_M = \Pi \oplus \chi$ and $\tau_M = \Pi \oplus L^1$ induce the vectorbundle morphisms

$$\begin{array}{rccc} P_{L^1}: & \Pi \oplus L^1 & \longrightarrow & \Pi \\ & X = X_\Pi + X_{L^1} & \longmapsto & X_\Pi \end{array} \qquad \begin{array}{rccc} P_\chi: & \Pi \oplus \chi & \longrightarrow & \Pi \\ & Y = Y_\Pi + Y_\chi & \longmapsto & Y_\Pi \end{array}$$

$$\begin{array}{rccc} P^{L^1}: & \Pi \oplus L^1 & \longrightarrow & L^1 \\ & X = X_\Pi + X_{L^1} & \longmapsto & X_{L^1} \end{array} \qquad \begin{array}{rccc} P^\chi: & \Pi \oplus \chi & \longrightarrow & \chi \\ & Y = Y_\Pi + Y_\chi & \longmapsto & Y_\chi \end{array}$$

So we get the morphisms

$$P_{L^1} : \mathcal{D}_1(M) \longrightarrow C^\infty(\Pi) \qquad \text{and} \qquad P_\chi : \mathcal{D}_1(M) \longrightarrow C^\infty(\Pi),$$

$$P^{L^1} : \mathcal{D}_1(M) \longrightarrow C^\infty\left(L^1\right) \qquad \text{and} \qquad P^\chi : \mathcal{D}_1(M) \longrightarrow C^\infty(\chi)$$

of $C^\infty(M)$-modules.

Remark 3.16 *We can represent the projections P_χ and P^χ as the tensors $P_\chi = 1 - X_1 \otimes \omega$ and $P^\chi = X_1 \otimes \omega$.*

$$\begin{aligned} P_\chi(\xi) &= \xi - \omega(\xi) X_1, \\ P^\chi(\xi) &= \omega(\xi) X_1. \end{aligned}$$

In the following we need to extend $j \in \Pi^* \otimes \Pi$. We have several possibilities. Two of them are more natural then the others: We can extend by zero along the characteristic vector field X_1 or along L^1. We choose to extend along the characteristic vector field.

Definition 3.17 *Define $\widetilde{j} \in T^*M \otimes TM$ in the following way:*

$$\widetilde{j} = \begin{cases} j & \text{on } \Pi \\ 0 & \text{on } \chi \end{cases}. \tag{3.6}$$

So $\widetilde{j}(X) = \widetilde{j}(X_\Pi + X_\chi) = j(X_\Pi)$. We notice that $P_\chi \circ \widetilde{j}(X) = j(X_\Pi) = j \circ P_\chi(X)$, hence

$$P_\chi \circ \widetilde{j} = j \circ P_\chi. \tag{3.7}$$

Definition 3.18 *Let $\xi, \eta \in C^\infty(\Pi)$. We define the **Nijenhuis tensor** to be:*

$$N_j(\xi, \eta) = P_\chi\left([j\xi, j\eta] - \widetilde{j}[j\xi, \eta] - \widetilde{j}[\xi, j\eta] + [\xi, \eta]\right). \tag{3.8}$$

Remark 3.19 *It is not clear why we should extend j along χ and project with P_χ in the definition of N_j. It seems like we could equally well extend j along L^1 and then define N_j by the projection P_{L^1} instead of P_χ. This is not an option however, since the Nijenhuis tensor will turn out to be identically zero.*

Lemma 3.20 *N_j is a tensor:*

$$N_j \in \Lambda^2 \Pi^* \otimes \Pi. \tag{3.9}$$

Proof. The proof is presented in appendix A. ■

Proposition 3.21 *The Nijenhuis tensor has the following properties:*

i) $N_j(X,Y) = -N_j(Y,X)$, *hence* $N_j(X,X) = 0$.

ii) $N_j(kX,Y) = kN_j(X,Y)$; $k \in \mathbb{R}$.

iii) $N_j(X_1 + X_2, Y) = N_j(X_1,Y) + N_j(X_2,Y)$.

iv) $N_j(jX,Y) = N_j(X,jY) = -jN_j(X,Y)$.

Here X and Y are vectors in Π, but these properties also hold for sections in Π.

Proof. The first 3 properties - skew symmetry and bilinearity - follows directly from the definition. We prove only the last property.

Let $X, Y \in \Pi$. First we calculate $N_j(jX,Y)$:

$$\begin{aligned}
N_j(jX,Y) &= P_\chi\left(\left[j^2X, jY\right] - \widetilde{j}\left[j^2X, Y\right] - \widetilde{j}\left[jX, jY\right] + \left[jX, Y\right]\right) \\
&= P_\chi\left(\left[X, jY\right] - \widetilde{j}\left[X, Y\right] - \widetilde{j}\left[jX, jY\right] + \left[jX, Y\right]\right) \\
&= P_\chi\left(\left[X, jY\right]_\Pi - j\left[X, Y\right]_\Pi - j\left[jX, jY\right]_\Pi + \left[jX, Y\right]_\Pi + \left[X, jY\right]_\chi + \left[jX, Y\right]_\chi\right) \\
&= P_\chi\left(\left[X, jY\right]_\Pi - j\left[X, Y\right]_\Pi - j\left[jX, jY\right]_\Pi + \left[jX, Y\right]_\Pi\right).
\end{aligned}$$

Second we calculate $N_j(X,jY)$:

$$\begin{aligned}
N_j(X,jY) &= P_\chi\left(\left[jX, j^2Y\right] - \widetilde{j}\left[jX, jY\right] - \widetilde{j}\left[X, j^2Y\right] + \left[X, jY\right]\right) \\
&= P_\chi\left(\left[jX, Y\right] - \widetilde{j}\left[jX, jY\right] - \widetilde{j}\left[X, Y\right] + \left[X, jY\right]\right) \\
&= N_j(jX,Y).
\end{aligned}$$

Finally we calculate $-jN_j(X,Y)$:

$$\begin{aligned}
-jN_j(X,Y) &= -j\circ P_\chi\left([jX,jY]-\widetilde{j}[jX,Y]-\widetilde{j}[X,jY]+[X,Y]\right)\\
&= -P_\chi\circ\widetilde{j}\left([jX,jY]-\widetilde{j}[jX,Y]-\widetilde{j}[X,jY]+[X,Y]\right)\\
&= -P_\chi\left(\widetilde{j}[jX,jY]-\widetilde{j^2}[jX,Y]-\widetilde{j^2}[X,jY]+\widetilde{j}[X,Y]\right)\\
&= -P_\chi\left(j[jX,jY]_\Pi-j^2[jX,Y]_\Pi-j^2[X,jY]_\Pi+j[X,Y]_\Pi\right)\\
&= -P_\chi\left(j[jX,jY]_\Pi-[jX,Y]_\Pi-[X,jY]_\Pi+j[X,Y]_\Pi\right)\\
&= P_\chi\left(-j[jX,jY]_\Pi+[jX,Y]_\Pi+[X,jY]_\Pi-j[X,Y]_\Pi\right)\\
&= N_j(jX,Y).
\end{aligned}$$

■

These properties coincide with the properties of the Nijenhuis tensor for the 4-dimensional symplectic case described in [Kr2], hence the name is justified. The proposition leads to the following corollaries:

Corollary 3.22 $N_j\left(\Pi^2_\pm,\Pi^2_\mp\right)=0.$

Proof. Let $X\in\Pi^2_+$ and $Y\in\Pi^2_-$. Then $jX=X$ and $jY=-Y$. From proposition 3.21 we get:

$$N_j(X,Y)=N_j(jX,Y)=N_j(X,jY)=N_j(X,-Y)=-N_j(X,Y).$$

So we see that $N_j\left(\Pi^2_\pm,\Pi^2_\mp\right)=0$. ■

Corollary 3.23 $N_j\left(\Pi^2_+,\Pi^2_+\right)\subseteq\Pi^2_-.$

Proof. Let $X\in\Pi^2_+$ and $Y\in\Pi^2_+$. Then $jX=X$ and $jY=Y$. From proposition 3.21 we get:

$$jN_j(X,Y)=-N_j(jX,Y)=-N_j(X,Y).$$

This implies that $N_j(X,Y)\in\Pi^2_-$. ■

Corollary 3.24 $N_j\left(\Pi^2_-,\Pi^2_-\right)\subseteq\Pi^2_+.$

Proof. Let $X\in\Pi^2_-$ and $Y\in\Pi^2_-$. Then $jX=-X$ and $jY=-Y$. From proposition 3.21 we get:

$$jN_j(X,Y)=-N_j(jX,Y)=-N_j(-X,Y)=N_j(X,Y).$$

This implies that $N_j(X,Y)\in\Pi^2_+$. ■

One can easily check that if ξ_+, η_+ and ξ_-, η_- generate Π^2_+ and Π^2_- correspondingly, then the following formulas hold:

$$\begin{aligned} N_j\left(\xi_+, \eta_+\right) &= 4P_- \circ P_\chi\left[\xi_+, \eta_+\right], \\ N_j\left(\xi_-, \eta_-\right) &= 4P_+ \circ P_\chi\left[\xi_-, \eta_-\right]. \end{aligned}$$

So we can write the Nijenhuis tensor like

$$N_j(\xi, \eta) = \begin{cases} 4P_\mp \circ P_\chi[\xi, \eta] & \text{when } \xi, \eta \in C^\infty\left(\Pi^2_\pm\right), \\ 0 & \text{when } \xi, \eta \in C^\infty\left(\Pi^2_\pm\right) \times C^\infty\left(\Pi^2_\mp\right). \end{cases} \tag{3.10}$$

Lemma 3.25 *The Nijenhuis tensor is identical to zero if and only if the vectorbundles χ and L^1 coincide.*

Proof. Let ξ_+, η_+ and ξ_-, η_- be local generating sections in Π^2_+ and Π^2_- correspondingly and let ℓ be a generating section in L^1. Then the characteristic vector field X_1 can be decomposed into the given generators as

$$X_1 = a_1\xi_+ + a_2\eta_+ + a_3\xi_- + a_4\eta_- + a_5\ell,$$

where $a_i \in C^\infty(M)$. Assume that $\left[\xi_+, \eta_+\right] \in C^\infty\left(\Pi^2_+\right) \oplus C^\infty\left(L^1\right)$ decomposes into $\left[\xi_+, \eta_+\right] = A\left[\xi_+, \eta_+\right]_+ + B\ell$ where $A, B \in C^\infty(M)$. Then we get

$$\begin{aligned} N_j\left(\xi_+, \eta_+\right) &= 4P_- \circ P_\chi\left(A\left[\xi_+, \eta_+\right]_+ + B\ell\right) = 4BP_- \circ P_\chi(\ell) \\ &= 4BP_- \circ P_\chi\left(\frac{1}{a_5}X_1 - \frac{a_1}{a_5}\xi_+ - \frac{a_2}{a_5}\eta_+ - \frac{a_3}{a_5}\xi_- - \frac{a_4}{a_5}\eta_-\right) \\ &= 4BP_-\left(\frac{a_1}{a_5}\xi_+ - \frac{a_2}{a_5}\eta_+ - \frac{a_3}{a_5}\xi_- - \frac{a_4}{a_5}\eta_-\right) = -4B\left(\frac{a_3}{a_5}\xi_- + \frac{a_4}{a_5}\eta_-\right). \end{aligned}$$

Assume that $\left[\xi_-, \eta_-\right] \in C^\infty\left(\Pi^2_-\right) \oplus C^\infty\left(L^1\right)$ decomposes into $\left[\xi_-, \eta_-\right] = \alpha\left[\xi_-, \eta_-\right]_- + \beta\ell$, where $\alpha, \beta \in C^\infty(M)$.

$$\begin{aligned} N_j\left(\xi_-, \eta_-\right) &= 4P_+ \circ P_\chi\left(\alpha\left[\xi_-, \eta_-\right]_- + \beta\ell\right) = 4\beta P_+ \circ P_\chi(\ell) \\ &= 4\beta P_+ \circ P_\chi\left(\frac{1}{a_5}X_1 - \frac{a_1}{a_5}\xi_+ - \frac{a_2}{a_5}\eta_+ - \frac{a_3}{a_5}\xi_- - \frac{a_4}{a_5}\eta_-\right) \\ &= 4\beta P_+\left(-\frac{a_1}{a_5}\xi_+ - \frac{a_2}{a_5}\eta_+ - \frac{a_3}{a_5}\xi_- - \frac{a_4}{a_5}\eta_-\right) = -4\beta\left(\frac{a_1}{a_5}\xi_+ + \frac{a_2}{a_5}\eta_+\right). \end{aligned}$$

We know that both B and β must be nonzero since Π^2_+ and Π^2_- are nonintegrable. So we conclude that $N_j = 0 \Longleftrightarrow a_1, a_2, a_3, a_4 = 0$. ■

Example 3.26 (Linear wave 3) *We continue our study of $M = \mathbb{R}^5(q_1, q_2, u, p_1, p_2)$ with the contact form $\omega = du - p_1dq_1 - p_2dq_2$, the contact distribution $\Pi = \langle \mathcal{D}_{q_1}, \mathcal{D}_{q_2}, \partial_{p_1}, \partial_{p_2} \rangle$*

and the hyperbolic effective form $\theta = dq_1 \wedge dp_2 + dq_2 \wedge dp_1$. *We know that* $X_1 = \partial_u$ *and* $L^1 = \Pi^3_+ \cap \Pi^3_- = \operatorname{span}\{\partial_u\}$ *in this case, so we conclude by the preceding lemma that* $N_j \equiv 0$.

Example 3.27 (Nonlinear 2) *Consider the same contact manifold as described in the preceding example. We know that the contact form* $\omega = du - p_1 dq_1 - p_2 dq_2$ *and the hyperbolic effective form*

$$\theta = dq_1 \wedge dp_1 - dq_2 \wedge dp_2 + S dq_2 \wedge dp_1 - T dp_1 \wedge dp_2$$

correspond to

$$\begin{aligned}
\Pi^2_+ &= \langle \mathcal{D}_{q_1}, T\mathcal{D}_{q_2} - 2\partial_{p_1} - S\partial_{p_2} \rangle, \qquad \Pi^2_- = \langle S\mathcal{D}_{q_1} - 2\mathcal{D}_{q_2}, T\mathcal{D}_{q_2} - S\partial_{p_2} \rangle, \\
j &= \begin{bmatrix} 1 & S & 0 & T \\ 0 & -1 & -T & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & S & -1 \end{bmatrix}, \\
L^1 &= \left\langle \left(S\mathcal{D}_{q_1}(T) - T\mathcal{D}_{q_1}(S) - \mathcal{D}_{q_2}(T) + \frac{\partial S}{\partial p_2} \right) \mathcal{D}_{q_1} - \mathcal{D}_{q_1}(T)\mathcal{D}_{q_2} - 2\partial_u + \mathcal{D}_{q_1}(S)\partial_{p_2} \right\rangle.
\end{aligned}$$

Let us denote

$$\begin{aligned}
\xi_+ &= \mathcal{D}_{q_1}, & \eta_+ &= T\mathcal{D}_{q_2} - 2\partial_{p_1} - S\partial_{p_2}, \\
\xi_- &= S\mathcal{D}_{q_1} - 2\mathcal{D}_{q_2}, & \eta_- &= T\mathcal{D}_{q_2} - S\partial_{p_2}.
\end{aligned}$$

Since $X_1 = \partial_u$ *we can calculate the Nijenhuis tensor by equation 3.10:*

$$\begin{aligned}
N_j(\xi_+, \eta_+) &= 4P_- \circ P_\chi [\xi_+, \eta_+] \\
&= 4P_- \circ P_\chi (\mathcal{D}_{q_1}(T)\mathcal{D}_{q_2} + 2\partial_u - \mathcal{D}_{q_1}(S)\partial_{p_2}) \\
&= 4P_- (\mathcal{D}_{q_1}(T)\mathcal{D}_{q_2} - \mathcal{D}_{q_1}(S)\partial_{p_2}) \\
&= 2(T\mathcal{D}_{q_1}(S) - S\mathcal{D}_{q_1}(T))\mathcal{D}_{q_1} + 4\mathcal{D}_{q_1}(T)\mathcal{D}_{q_2} - 4\mathcal{D}_{q_1}(S)\partial_{p_2}.
\end{aligned}$$

Similarly one can check that

$$\begin{aligned}
N_j(\xi_-, \eta_-) &= 4P_+ \circ P_\chi [S\mathcal{D}_{q_1} - 2\mathcal{D}_{q_2}, T\mathcal{D}_{q_2} - S\partial_{p_2}] \\
&= 2S \left(S\mathcal{D}_{q_1}(T) - T\mathcal{D}_{q_1}(S) - 2\mathcal{D}_{q_2}(T) + 2\frac{\partial S}{\partial p_2} \right) \mathcal{D}_{q_1}.
\end{aligned}$$

It is clear that the Nijenhuis tensor defines a vectorbundle by its image. We will only study the cases when N_j defines a vectorbundle with constant rank. This is clearly not always the case.

Definition 3.28 *We define* $N_j(\Pi^2_\mp, \Pi^2_\mp) = R^1_\pm \subseteq \Pi^2_\pm$.

Definition 3.29 *We call* $\Pi_j = \mathrm{Im}\,(N_j) = R^1_+ \oplus R^1_-$ *the* ***characteristic distribution*** *of the Nijenhuis tensor.*

Obviously Π_j projects onto R^1_+ and R^1_- under the projections P_+ and P_- correspondingly. We also notice that $N_j\,(\Pi_j, \Pi_j) = 0$.

Proposition 3.30 *Any submanifold* N *of the strict contact manifold* M *satisfying* $TN = \Pi_j$ *is Legendrian.*

Proof. Obviously N is an integral manifold of Π and $\dim N = \mathrm{rank}\,\Pi_j = 2 = \frac{\dim M - 1}{2}$, hence N satisfies definition 2.15. ■

One should also note that $TN = \Pi_j$ requires that Π_j is integrable.

From the properties of N_j it follows that there can be 3 main cases:

1. The Nijenhuis tensor is identically zero, $N_j \equiv 0$.
2. Π_j is a rank 1 distribution within Π^2_+ or Π^2_-.
3. Π_j is a rank 2 distribution.

We will consider case 3. From corollary 3.8 we know that $\partial\Pi_j$ must be a subbundle of Π. Under the assumption that $\partial\Pi_j$ has constant rank there are 4 cases:

- $\partial\Pi_j = \Pi_j$, i.e. Π_j is integrable.
- $\partial\Pi_j \neq \Pi_j$ and $\partial\Pi_j \cap \Pi^2_+ = \Pi^2_+$, i.e. the commutator of the two generating sections in Π_j is a section in Π^2_+.
- $\partial\Pi_j \neq \Pi_j$ and $\partial\Pi_j \cap \Pi^2_+ = \Pi^2_-$, i.e. the commutator of the two generating sections in Π_j is a section in Π^2_-.
- $\partial\Pi_j \neq \Pi_j$ and $\partial\Pi_j \cap \Pi^2_+$, $\partial\Pi_j \cap \Pi^2_+$ are both 1-dimensional, i.e. the commutator of the two generating sections in Π_j is neither a section in Π^2_+, Π^2_- nor Π_j.

The following picture shows how the fibers $\Pi_j\,(x)\,, \Pi^2_+\,(x)\,, \Pi^2_-\,(x)\,, L^1\,(x)$ and $\chi\,(x)$ relate in T_xM when Π_j has rank 2:

Example 3.31 (Nonlinear 3) *We continue example 3.27. We see that*

$$\begin{aligned} R^1_+ &= \langle \mathcal{D}_{q_1} \rangle = \left\langle \xi_+ \right\rangle, \\ R^1_- &= \langle (T\mathcal{D}_{q_1}(S) - S\mathcal{D}_{q_1}(T))\,\mathcal{D}_{q_1} + 2\mathcal{D}_{q_1}(T)\,\mathcal{D}_{q_2} - 2\mathcal{D}_{q_1}(S)\,\partial_{p_2} \rangle = \left\langle \eta_- \right\rangle. \end{aligned}$$

Chapter 4

Invariants of hyperbolic M-A equations

We will now introduce three new invariants, the 1-form σ, the distribution Γ_σ^1 and the vectorvalued 2-form R_j^σ, of the generalized hyperbolic Monge-Ampere equation (ω, θ). In the first two sections we construct and investigate these invariants. Section 3 contains the Main Theorem, stating that any nondegenerate equation (ω, θ) defines a canonical frame on M.

4.1 The canonical 1-form σ and the distribution Γ_σ^1

In this section we construct a canonical 1-form $\sigma \in C^\infty(\Pi^*)$ together with a rank 1 distribution Γ_σ^1 from the forms $\omega \in \Omega^1(M)$ and $\Omega, i_j\Omega \in C^\infty(\Lambda^2\Pi^*)$. The construction is based on the properties of the de-Rham complex

$$C^\infty(M) \xrightarrow{d} \Omega^1(M) \xrightarrow{d} \Omega^2(M) \xrightarrow{d} \Omega^3(M) \xrightarrow{d} \Omega^4(M) \xrightarrow{d} \Omega^5(M),$$

where M is our 5-dimensional manifold.

The canonical splitting

$$\tau_M^* = \Pi^* \oplus \chi^*$$

gives us the following isomorphisms:

$$\begin{aligned}
\Omega^1(M) &\simeq C^\infty(\Pi^*) \oplus_{C^\infty(M)} C^\infty(\chi^*), \\
\Omega^2(M) &\simeq C^\infty\left(\Lambda^2\Pi^*\right) \oplus_{C^\infty(M)} C^\infty(\Pi^* \wedge \chi^*), \\
\Omega^3(M) &\simeq C^\infty\left(\Lambda^3\Pi^*\right) \oplus_{C^\infty(M)} C^\infty\left(\Lambda^2\Pi^* \wedge \chi^*\right), \\
\Omega^4(M) &\simeq C^\infty\left(\Lambda^4\Pi^*\right) \oplus_{C^\infty(M)} C^\infty\left(\Lambda^3\Pi^* \wedge \chi^*\right).
\end{aligned}$$

Define the following two $C^\infty(M)$-linear maps:

$$\begin{array}{ccccc} C^\infty(\Lambda^i\Pi^*) & \stackrel{\text{Ext}}{\longrightarrow} & \Omega^i(M) & \text{and} & \Omega^i(M) & \stackrel{\text{Rst}}{\longrightarrow} & C^\infty(\Lambda^i\Pi^*) \\ \mu & \longmapsto & \widetilde{\mu} & & \omega & \longmapsto & \omega|_\Pi = \omega \bmod \chi^* \end{array}, \tag{4.1}$$

where

$$\widetilde{\mu} = \begin{cases} \mu & \text{on } \Lambda^i\Pi \\ 0 & \text{on } \Lambda^{i-1}\Pi \wedge \chi \end{cases}.$$

Definition 4.1 *Define the operator d_Π in the following way:*

$$d_\Pi = \begin{cases} \text{Rst} \circ d \circ \text{Ext} & ; \text{ on } C^\infty(\Lambda^i\Pi^*); i \geq 1, \\ \text{Rst} \circ d & ; \text{ on } C^\infty(M). \end{cases}$$

Then, for $i = 0, 1, 2, 3$ we have the following commutative diagram:

$$\begin{array}{ccc} \Omega^i(M) & \xrightarrow{d} & \Omega^{i+1}(M) \\ \uparrow{\scriptstyle Ext} & & \downarrow{\scriptstyle Rst} \\ C^\infty(\Lambda^i\Pi^*) & \xrightarrow{d_\Pi} & C^\infty(\Lambda^{i+1}\Pi^*) \end{array}$$

From the effective form θ we get the canonical 3-form $d_\Pi\theta \in C^\infty(\Lambda^3\Pi^*)$. From [Ly1] and [Ly2] we know that

$$\begin{array}{rccc} \top: & C^\infty(\Pi^*) & \longrightarrow & C^\infty(\Lambda^3\Pi^*) \\ & \alpha & \longmapsto & \alpha \wedge \Omega \end{array}$$

is an isomorphism. Hence there exists an unique 1-form $\sigma \in C^\infty(\Pi^*)$ such that $\top(\sigma) = d_\Pi\theta$.

Definition 4.2 *The canonical 1-form $\sigma \in C^\infty(\Pi^*)$ is defined to be the preimage of $d_\Pi\theta$ under the isomorphism $\top$.*

This 1-form induces a section in Π.

Definition 4.3 *We define the vector field $X_\sigma \in C^\infty(\Pi)$ by:*

$$i_{X_\sigma}\Omega = \sigma,$$

where Ω is the symplectic structure on Π.

Obviously X_σ is uniquely defined by σ since Ω is nondegenerate. We notice that

$$\sigma(X_\sigma) = 0.$$

Definition 4.4 *Define the distribution* Γ_σ^1 *to be*

$$\Gamma_\sigma^1 = \langle X_\sigma \rangle .$$

Example 4.5 *Consider* $M = \mathbb{R}^5 (q_1, q_2, u, p_1, p_2)$ *with the contact distribution* $\Pi = \langle \mathcal{D}_{q_1}, \mathcal{D}_{q_2}, \partial_{p_1}, \partial_{p_2} \rangle$*, the symplectic structure* $\Omega = dq_1 \wedge dp_1 + dq_2 \wedge dp_2$ *and the general effective form*

$$\theta = F_1 dq_1 \wedge dq_2 + F_2 dq_2 \wedge dp_1 + F_3 dq_1 \wedge dp_2 + F_4 dq_1 \wedge dp_1 - F_4 dq_2 \wedge dp_2 + F_5 dp_1 \wedge dp_2,$$

where $F_1, \ldots, F_5 \in C^\infty(M)$*. We know that* $X_1 = \partial_u$*. Let us identify*

$$\begin{aligned} \chi^* &\simeq \operatorname{Ann}(\Pi) = \langle du - p_1 dq_1 - p_2 dq_2 \rangle , \\ \Pi^* &\simeq \operatorname{Ann}(\chi) = \langle dq_1, dq_2, dp_1, dp_2 \rangle . \end{aligned}$$

Then one can check that

$$\Lambda^3 \Pi^* = \langle dq_1 \wedge dq_2 \wedge dp_1, dq_1 \wedge dq_2 \wedge dp_2, dq_1 \wedge dp_1 \wedge dp_2, dq_2 \wedge dp_1 \wedge dp_2 \rangle ,$$

hence

$$\begin{aligned} d_\Pi \theta &= \left(\frac{\partial F_1}{\partial p_1} - \mathcal{D}_{q_2}(F_4) + \mathcal{D}_{q_1}(F_2) \right) dq_1 \wedge dq_2 \wedge dp_1 \\ &\quad + \left(\frac{\partial F_1}{\partial p_2} - \mathcal{D}_{q_2}(F_3) - \mathcal{D}_{q_1}(F_4) \right) dq_1 \wedge dq_2 \wedge dp_2 \\ &\quad + \left(\frac{\partial F_4}{\partial p_2} - \frac{\partial F_3}{\partial p_1} + \mathcal{D}_{q_1}(F_5) \right) dq_1 \wedge dp_1 \wedge dp_2 \\ &\quad + \left(\frac{\partial F_2}{\partial p_2} + \frac{\partial F_4}{\partial p_1} + \mathcal{D}_{q_2}(F_5) \right) dq_2 \wedge dp_1 \wedge dp_2 . \end{aligned}$$

Let $\sigma = \lambda_1 dq_1 + \lambda_2 dq_2 + \lambda_3 dp_1 + \lambda_4 dp_2 \in \Pi^*$*. Then*

$$\sigma \wedge \Omega = -\lambda_2 dq_1 \wedge dq_2 \wedge dp_1 + \lambda_1 dq_1 \wedge dq_2 \wedge dp_2 + \lambda_4 dq_1 \wedge dp_1 \wedge dp_2 - \lambda_3 dq_2 \wedge dp_1 \wedge dp_2 .$$

By comparing $d_\Pi \theta$ *with* $\sigma \wedge \Omega$ *we conclude that*

$$\begin{aligned} \sigma &= \left(\frac{\partial F_1}{\partial p_2} - \mathcal{D}_{q_2}(F_3) - \mathcal{D}_{q_1}(F_4) \right) dq_1 - \left(\frac{\partial F_1}{\partial p_1} - \mathcal{D}_{q_2}(F_4) + \mathcal{D}_{q_1}(F_2) \right) dq_2 \\ &\quad - \left(\frac{\partial F_2}{\partial p_2} + \frac{\partial F_4}{\partial p_1} + \mathcal{D}_{q_2}(F_5) \right) dp_1 + \left(\frac{\partial F_4}{\partial p_2} - \frac{\partial F_3}{\partial p_1} + \mathcal{D}_{q_1}(F_5) \right) dp_2 . \end{aligned}$$

This leads to

$$\begin{aligned} X_\sigma &= -\left(\frac{\partial F_2}{\partial p_2}+\frac{\partial F_4}{\partial p_1}+\mathcal{D}_{q_2}(F_5)\right)\mathcal{D}_{q_1}+\left(\frac{\partial F_4}{\partial p_2}-\frac{\partial F_3}{\partial p_1}+\mathcal{D}_{q_1}(F_5)\right)\mathcal{D}_{q_2} \\ &\quad -\left(\frac{\partial F_1}{\partial p_2}-\mathcal{D}_{q_2}(F_3)-\mathcal{D}_{q_1}(F_4)\right)\partial_{p_1}+\left(\frac{\partial F_1}{\partial p_1}-\mathcal{D}_{q_2}(F_4)+\mathcal{D}_{q_1}(F_2)\right)\partial_{p_2}. \end{aligned}$$

In the case when θ has constant coefficients, e.g. in the case

$$\frac{\partial^2 h}{\partial q_1^2}-\frac{\partial^2 h}{\partial q_2^2}=0 \qquad \Longleftrightarrow \qquad \theta = dp_1\wedge dq_2+dp_2\wedge dq_1,$$

we see that $\sigma = 0$. While in the case

$$-S\frac{\partial^2 u}{\partial q_1^2}+2\frac{\partial^2 u}{\partial q_1\partial q_2}-T\left(\frac{\partial^2 u}{\partial q_1^2}\frac{\partial^2 u}{\partial q_2^2}-\left(\frac{\partial^2 u}{\partial q_1\partial q_2}\right)^2\right)=0,$$

$$\theta = dq_1\wedge dp_1-dq_2\wedge dp_2+Sdq_2\wedge dp_1-Tdp_1\wedge dp_2,$$

we find that $\sigma = -\mathcal{D}_{q_1}(S)\,dq_2+\left(\mathcal{D}_{q_2}(T)-\frac{\partial S}{\partial p_2}\right)dp_1-\mathcal{D}_{q_1}(T)\,dp_2$.

4.2 The vectorvalued 2-form R_j^σ

In this section we define a vectorvalued 2-form $R_j^\sigma \in \Lambda^2\Pi^*\otimes\Pi$ from the 1-form σ and the Nijenhuis tensor. The properties of R_j^σ contain some useful information concerning the distribution Γ_σ^1. When investigating R_j^σ, we need to extend θ and Ω to $\widetilde{\theta},\widetilde{\Omega}\in\Omega^2(M)$ by the extension defined in equation 4.1. We can observe that

$$\widetilde{\Omega}=\left\{\begin{array}{lll} d\omega & ; & \text{on } \Pi \\ 0 & ; & \text{on } \chi \end{array}\right. = d\omega.$$

The last equality follows from the definition of the characteristic vector field X_1 that generates χ.

Definition 4.6 *Let $\xi,\eta\in C^\infty(\Pi)$. We define the vector valued 2-form $R_j^\sigma\in\Lambda^2\Pi^*\otimes\Pi$ to be the following:*

$$R_j^\sigma(\xi,\eta)=N_j(\xi,\eta)+j\eta\sigma(\xi)-j\xi\sigma(\eta)+\eta\sigma(j\xi)-\xi\sigma(j\eta). \tag{4.2}$$

Proposition 4.7 *$R_j^\sigma(\xi,\eta)$ has the following properties:*

i) $R_j^\sigma(\xi,\eta)=-R_j^\sigma(\eta,\xi)$.

ii) $R_j^\sigma(k\xi,\eta)=kR_j^\sigma(\xi,\eta)$; $k\in\mathbb{R}$.

iii) $R_j^\sigma(\xi_1+\xi_2,\eta)=R_j^\sigma(\xi_1,\eta)+R_j^\sigma(\xi_2,\eta)$.

iv) $R_j^\sigma(j\xi,\eta)=R_j^\sigma(\xi,j\eta)$.

v) $2N_j(j\xi,\eta)=R_j^\sigma(j\xi,\eta)-jR_j^\sigma(\xi,\eta)$.

Proof. All five properties follow directly from the properties of the Nijenhuis tensor. We show only the last one.

$$\begin{aligned}
2N_j(j\xi,\eta) &= N_j(j\xi,\eta)+N_j(j\xi,\eta)=N_j(j\xi,\eta)-jN_j(\xi,\eta)\\
&= R_j^\sigma(j\xi,\eta)-j\eta\sigma(j\xi)+\xi\sigma(\eta)-\eta\sigma(\xi)+j\xi\sigma(j\eta)\\
&\quad -j\left(R_j^\sigma(\xi,\eta)-j\eta\sigma(\xi)+j\xi\sigma(\eta)-\eta\sigma(j\xi)+\xi\sigma(j\eta)\right)\\
&= R_j^\sigma(j\xi,\eta)-j\eta\sigma(j\xi)+\xi\sigma(\eta)-\eta\sigma(\xi)+j\xi\sigma(j\eta)\\
&\quad -jR_j^\sigma(\xi,\eta)+\eta\sigma(\xi)-\xi\sigma(\eta)+j\eta\sigma(j\xi)-j\xi\sigma(j\eta)\\
&= R_j^\sigma(j\xi,\eta)-jR_j^\sigma(\xi,\eta).
\end{aligned}$$

■

Theorem 4.8 *The formula*

$$R_j^\sigma=-2jX_\sigma\otimes\Omega \tag{4.3}$$

holds.

To prove this theorem we need some results.

Proposition 4.9 *Let $\Omega\in C^\infty(\Lambda^2\Pi^*)$ be the symplectic form and $\theta=i_j\Omega$ the effective form. Then the following formula holds:*

$$\begin{aligned}
d\widetilde{\theta}(\xi,\eta,\gamma) &= L_\xi\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right)-L_{\widetilde{j}\xi}\left(\widetilde{\Omega}(\eta,\gamma)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right)\\
&\quad -\widetilde{\Omega}\left(\widetilde{j}[\xi,\eta],\gamma\right)+\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right)-\widetilde{\Omega}\left(\eta,\widetilde{j}[\xi,\gamma]\right)\\
&\quad +d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right)
\end{aligned} \tag{4.4}$$

for vector fields $\xi,\eta,\gamma\in\mathcal{D}_1(M)$.

Proof. The proof is presented in appendix B. ■

We know that the vectorbundle isomorphism $\tau_M\simeq\Pi\oplus\chi$ induces the $C^\infty(M)$-linear isomorphism

$$\mathcal{D}_1(M)\simeq C^\infty(\Pi)\oplus_{C^\infty(M)}C^\infty(\chi).$$

Restricting the proposition to $C^\infty(\Pi)$ we obtain the following corollary.

Corollary 4.10 *The formula*

$$\begin{aligned} d_{\Pi}\theta\left(\xi,\eta,\gamma\right) &= L_{\xi}\left(\widetilde{\Omega}\left(j\eta,\gamma\right)\right)-L_{j\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[j\xi,\eta\right],\gamma\right) \\ &\quad -\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)+\widetilde{\Omega}\left(\eta,\left[j\xi,\gamma\right]\right)-\widetilde{\Omega}\left(\eta,\widetilde{j}\left[\xi,\gamma\right]\right) \end{aligned} \tag{4.5}$$

holds for $\xi,\eta,\gamma\in C^{\infty}\left(\Pi\right)$.

Proof. From the proposition we have:

$$\begin{aligned} d\widetilde{\theta}\left(\xi,\eta,\gamma\right) &= L_{\xi}\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right)-L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right)-\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right) \\ &\quad +\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right)-\widetilde{\Omega}\left(\eta,\widetilde{j}\left[\xi,\gamma\right]\right)+d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right). \end{aligned}$$

Since $\xi,\eta,\gamma\in C^{\infty}\left(\Pi\right)$, it follows from the definition of d_{Π} that $d\widetilde{\theta}\left(\xi,\eta,\gamma\right)=d_{\Pi}\theta\left(\xi,\eta,\gamma\right)$. We also know that $d\widetilde{\Omega}\left(\xi,\widetilde{j}\eta,\gamma\right)=d\left(d\omega\right)\left(\xi,\widetilde{j}\eta,\gamma\right)=0$, so we get:

$$\begin{aligned} d_{\Pi}\theta\left(\xi,\eta,\gamma\right) &= L_{\xi}\left(\widetilde{\Omega}\left(j\eta,\gamma\right)\right)-L_{j\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[j\xi,\eta\right],\gamma\right) \\ &\quad -\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)+\widetilde{\Omega}\left(\eta,\left[j\xi,\gamma\right]\right)-\widetilde{\Omega}\left(\eta,\widetilde{j}\left[\xi,\gamma\right]\right). \end{aligned}$$

■

Lemma 4.11 *Let* $\xi,\eta\in C^{\infty}\left(\Pi\right)$. *Then*

$$i_{N_{j}(\xi,\eta)}\theta=-d_{\Pi}\theta\left(\xi,\eta,\bullet\right)-d_{\Pi}\theta\left(j\xi,j\eta,\bullet\right). \tag{4.6}$$

Proof. Let $\xi,\eta,\gamma\in C^{\infty}\left(\Pi\right)\subset\mathcal{D}_{1}\left(M\right)$. From the preceding corollary we know that

$$\begin{aligned} -d_{\Pi}\theta\left(\xi,\eta,\gamma\right) &= -L_{\xi}\left(\widetilde{\Omega}\left(j\eta,\gamma\right)\right)+L_{j\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)-\widetilde{\Omega}\left(\left[j\xi,\eta\right],\gamma\right) \\ &\quad +\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)-\widetilde{\Omega}\left(\eta,\left[j\xi,\gamma\right]\right)+\widetilde{\Omega}\left(\eta,\widetilde{j}\left[\xi,\gamma\right]\right). \end{aligned}$$

Since $j^{2}=1$ we also get:

$$\begin{aligned} -d_{\Pi}\theta\left(j\xi,j\eta,\gamma\right) &= -L_{j\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+L_{\xi}\left(\widetilde{\Omega}\left(j\eta,\gamma\right)\right)-\widetilde{\Omega}\left(\left[\xi,j\eta\right],\gamma\right) \\ &\quad +\widetilde{\Omega}\left(\widetilde{j}\left[j\xi,j\eta\right],\gamma\right)-\widetilde{\Omega}\left(j\eta,\left[\xi,\gamma\right]\right)+\widetilde{\Omega}\left(j\eta,\widetilde{j}\left[j\xi,\gamma\right]\right). \end{aligned}$$

Considering their sum, we see:

$$\begin{aligned}-d_{\Pi}\theta(\xi,\eta,\gamma)-d_{\Pi}\theta(j\xi,j\eta,\gamma) &= \widetilde{\Omega}\left(\widetilde{j}[j\xi,j\eta],\gamma\right)-\widetilde{\Omega}([j\xi,\eta],\gamma)+\widetilde{\Omega}\left(\widetilde{j}[\xi,\eta],\gamma\right)\\ &\quad -\widetilde{\Omega}([\xi,j\eta],\gamma)-\widetilde{\Omega}(\eta,[j\xi,\gamma])+\widetilde{\Omega}\left(j\eta,\widetilde{j}[j\xi,\gamma]\right)\\ &\quad +\widetilde{\Omega}\left(\eta,\widetilde{j}[\xi,\gamma]\right)-\widetilde{\Omega}(j\eta,[\xi,\gamma]).\end{aligned}$$

One can easily check that $\widetilde{\Omega}\left(\widetilde{j}X,Y\right)=\widetilde{\Omega}\left(X,\widetilde{j}Y\right)$ and since $\widetilde{j}j\equiv 1 \bmod \chi$, the last four terms vanish. So we are left with:

$$\begin{aligned}-d_{\Pi}\theta(\xi,\eta,\gamma)-d_{\Pi}\theta(j\xi,j\eta,\gamma) &= \widetilde{\Omega}\left(\widetilde{j}[j\xi,j\eta],\gamma\right)-\widetilde{\Omega}([j\xi,\eta],\gamma)-\widetilde{\Omega}([\xi,j\eta],\gamma)\\ &\quad +\widetilde{\Omega}\left(\widetilde{j}[\xi,\eta],\gamma\right)\\ &= \widetilde{\Omega}\left(\widetilde{j}[j\xi,j\eta]-[j\xi,\eta]-[\xi,j\eta]+\widetilde{j}[\xi,\eta],\gamma\right)\\ &= \widetilde{\theta}\left([j\xi,j\eta]-\widetilde{j}[j\xi,\eta]-\widetilde{j}[\xi,j\eta]+[\xi,\eta],\gamma\right),\end{aligned}$$

where we have used that $i_j\Omega=\theta \Longleftrightarrow i_{\widetilde{j}}\widetilde{\Omega}=\widetilde{\theta}$. Since $\widetilde{\theta}=0$ on χ we can write:

$$\begin{aligned}-d_{\Pi}\theta(\xi,\eta,\gamma)-d_{\Pi}\theta(j\xi,j\eta,\gamma) &= \theta\left(P_{\chi}\left([j\xi,j\eta]-\widetilde{j}[j\xi,\eta]-\widetilde{j}[\xi,j\eta]+[\xi,\eta]\right),\gamma\right)\\ &= \theta(N_j(\xi,\eta),\gamma).\end{aligned}$$

The claim follows. ■

Lemma 4.12 *Let $\xi,\eta,\gamma\in C^{\infty}(\Pi)$. Then the following identity holds:*

$$d_{\Pi}\theta(\xi,\eta,\gamma)=\Omega(\xi,\eta)\sigma(\gamma)+\Omega(\gamma,\xi)\sigma(\eta)+\Omega(\eta,\gamma)\sigma(\xi). \tag{4.7}$$

Proof. We know from the construction of σ that

$$d_{\Pi}\theta=\Omega\wedge\sigma.$$

Let $\xi,\eta,\gamma\in C^{\infty}(\Pi)$. Simple calculation gives the wanted result.

$$\begin{aligned}d_{\Pi}\theta(\xi,\eta,\gamma) &= i_{\gamma}i_{\eta}i_{\xi}(\Omega\wedge\sigma)=i_{\gamma}i_{\eta}(\Omega(\xi,\bullet)\wedge\sigma+\Omega\sigma(\xi))\\ &= i_{\gamma}(\Omega(\xi,\eta)\sigma-\Omega(\xi,\bullet)\sigma(\eta)+\Omega(\eta,\bullet)\sigma(\xi))\\ &= \Omega(\xi,\eta)\sigma(\gamma)-\Omega(\xi,\gamma)\sigma(\eta)+\Omega(\eta,\gamma)\sigma(\xi)\\ &= \Omega(\xi,\eta)\sigma(\gamma)+\Omega(\gamma,\xi)\sigma(\eta)+\Omega(\eta,\gamma)\sigma(\xi).\end{aligned}$$

■

Now we have what we need to prove **theorem 4.8**:

Proof. Let $\xi, \eta, \gamma \in C^\infty(\Pi)$. From lemma 4.11 and 4.12 it follows that

$$\begin{aligned}
\Omega(jN_j(\xi,\eta),\gamma) &= \theta(N_j(\xi,\eta),\gamma) = -d_\Pi\theta(\xi,\eta,\gamma) - d_\Pi\theta(j\xi,j\eta,\gamma) \\
&= -\Omega(\xi,\eta)\sigma(\gamma) - \Omega(\gamma,\xi)\sigma(\eta) - \Omega(\eta,\gamma)\sigma(\xi) - \Omega(j\xi,j\eta)\sigma(\gamma) \\
&\quad -\Omega(\gamma,j\xi)\sigma(j\eta) - \Omega(j\eta,\gamma)\sigma(j\xi) \\
&= -2\Omega(\xi,\eta)\sigma(\gamma) - \Omega(\gamma,\xi)\sigma(\eta) - \Omega(\gamma,j\xi)\sigma(j\eta) - \Omega(\eta,\gamma)\sigma(\xi) \\
&\quad -\Omega(j\eta,\gamma)\sigma(j\xi).
\end{aligned}$$

Thus

$$\begin{aligned}
-2\Omega(\xi,\eta)\sigma(\gamma) &= \Omega(jN_j(\xi,\eta),\gamma) + \Omega(\gamma,\xi)\sigma(\eta) + \Omega(\gamma,j\xi)\sigma(j\eta) + \Omega(\eta,\gamma)\sigma(\xi) \\
&\quad +\Omega(j\eta,\gamma)\sigma(j\xi) \\
&= \Omega(jN_j(\xi,\eta),\gamma) - \Omega(\sigma(\eta)\xi,\gamma) - \Omega(\sigma(j\eta)j\xi,\gamma) + \Omega(\sigma(\xi)\eta,\gamma) \\
&\quad +\Omega(\sigma(j\xi)j\eta,\gamma) \\
&= \Omega(jN_j(\xi,\eta) - \sigma(\eta)\xi - \sigma(j\eta)j\xi + \sigma(\xi)\eta + \sigma(j\xi)j\eta,\gamma) \\
&= \Omega\left(jR_j^\sigma(\xi,\eta),\gamma\right).
\end{aligned}$$

From this equation we find:

$$\begin{aligned}
\Omega\left(jR_j^\sigma(\xi,\eta),\gamma\right) &= -2\widetilde{\Omega}(\xi,\eta)\sigma(\gamma) = \sigma(-2\Omega(\xi,\eta)\cdot\gamma) \\
&= \Omega\left(X_\sigma, -2\widetilde{\Omega}(\xi,\eta)\cdot\gamma\right) \\
&= \Omega\left(-2\widetilde{\Omega}(\xi,\eta)\cdot X_\sigma,\gamma\right).
\end{aligned}$$

Since Ω is nondegenerate, this implies that

$$jR_j^\sigma(\xi,\eta) = -2\Omega(\xi,\eta)\cdot X_\sigma,$$

and the claim follows. ■

Remark 4.13 *We observe the following chain of relations:*

$$d_\Pi\theta = 0 \overset{1}{\Longleftrightarrow} \sigma = 0 \overset{2}{\Longleftrightarrow} X_\sigma = 0 \overset{3}{\Longleftrightarrow} R_j^\sigma = 0 \overset{4}{\Longleftrightarrow} N_j = 0.$$

The first equivalence follows from the construction of σ. The second follows from the construction of X_σ. The third one follows from theorem 4.8, since Ω is nondegenerate. And the fourth equivalence follows from proposition 4.7 v).

Lemma 4.14 $\Gamma_\sigma^1 \subset \Pi_j \subset \ker(\sigma) = (\Gamma_\sigma^1)^{\perp\Omega}$.

Proof. By the definition of X_σ, we have that $\Omega(X_\sigma,\xi) = \sigma(\xi)$. So $\ker(\sigma) = (\Gamma_\sigma^1)^{\perp\Omega}$.

Obviously $\sigma(X_\sigma) = 0$. From this it follows that

$$\begin{aligned}\sigma(N_j(\xi,\eta)) &= \sigma(jN_j(j\eta,\xi)) = \sigma\left(jR_j^\sigma(j\eta,\xi) + \xi\sigma(j\eta) - j\eta\sigma(\xi) + j\xi\sigma(\eta) - \eta\sigma(j\xi)\right)\\ &= \sigma\left(jR_j^\sigma(j\eta,\xi)\right) + \sigma(j\eta)\sigma(\xi) - \sigma(\xi)\sigma(j\eta) + \sigma(\eta)\sigma(j\xi) - \sigma(j\xi)\sigma(\eta)\\ &= \sigma\left(jR_j^\sigma(j\eta,\xi)\right) = \sigma(-2\Omega(j\eta,\xi)X_\sigma) = 0.\end{aligned}$$

Thus $\Pi_j \subset \ker(\sigma)$.

Let us show that $\Gamma_\sigma^1 \subset \Pi_j$. By construction, Π_j projects to R_+^1 and R_-^1, that is:

$$\begin{aligned}P_-(\Pi_j) &= R_-^1,\\ P_+(\Pi_j) &= R_+^1.\end{aligned}$$

Let X and Y be vectors in Π. From proposition 4.7 it follows that

$$\begin{aligned}-2jN_j(X,Y) &= R_j^\sigma(jX,Y) - jR_j^\sigma(X,Y),\\ 2N_j(X,Y) &= R_j^\sigma(X,Y) - jR_j^\sigma(jX,Y).\end{aligned}$$

By adding and subtracting these equations we get:

$$\begin{aligned}2(1-j)N_j(X,Y) &= (1-j)\left(R_j^\sigma(jX,Y) + R_j^\sigma(X,Y)\right) = (1-j)R_j^\sigma((1+j)X,Y),\\ 2(1+j)N_j(X,Y) &= (1+j)\left(R_j^\sigma(X,Y) - R_j^\sigma(jX,Y)\right) = (1+j)R_j^\sigma((1-j)X,Y).\end{aligned}$$

Using theorem 4.8, we see:

$$\begin{aligned}(1-j)R_j^\sigma((1+j)X,Y) &= -2\Omega((1+j)X,Y)(1-j)jX_\sigma = 2\Omega(2P_+X,Y)(1-j)X_\sigma,\\ (1+j)R_j^\sigma((1-j)X,Y) &= -2\Omega((1-j)X,Y)(1+j)jX_\sigma = -2\Omega(2P_-X,Y)(1+j)X_\sigma.\end{aligned}$$

Since Π_+^2 and Π_-^2 are skew orthogonal with respect to Ω, we can write

$$\begin{aligned}P_-N_j(X,Y) &= \Omega(P_+X,P_+Y)(1-j)X_\sigma,\\ P_+N_j(X,Y) &= -\Omega(P_-X,P_-Y)(1+j)X_\sigma.\end{aligned}$$

We see that $(1\pm j)X_\sigma \in R_\pm^1$. Since $X_\sigma = P_+X_\sigma + P_-X_\sigma$, we conclude that $X_\sigma \in R_+^1 \oplus R_-^1 = \Pi_j$. This implies that $\Gamma_\sigma^1 \subset \Pi_j$ as we were to prove. ■

Corollary 4.15 $\Gamma_\sigma^1 \cap R_+^1 = 0$ *and* $\Gamma_\sigma^1 \cap R_-^1 = 0$.

Proof. We know the equations

$$P_-N_j(X,Y) = \Omega(P_+X,P_+Y)(1-j)X_\sigma$$

and

$$P_+N_j(X,Y) = -\Omega(P_-X,P_-Y)(1+j)X_\sigma.$$

Assume now that $\Gamma^1_\sigma \cap R^1_+ \neq 0$, that is $\Gamma^1_\sigma = R^1_+$. Then $P_- X_\sigma = 0$ must be true, hence

$$P_- N_j(X, Y) = 0$$

must hold for all vectors $X, Y \in \Pi$. We know that $P_- N_j\left(\Pi^2_+, \Pi^2_+\right) = R^1_- \neq 0$, hence the assumption must be wrong and $\Gamma^1_\sigma \cap R^1_+ = 0$ must be true. Similarly we can prove that $\Gamma^1_\sigma \cap R^1_- = 0$. ■

Now we have two decompositions of Π_j, $R^1_+ \oplus R^1_-$ and $\Gamma^1_\sigma \oplus j\Gamma^1_\sigma$.

4.3 The canonical $\{e\}$-structure

Let ξ_+ and η_- be generating sections in R^1_+ and R^1_- in a neighborhood $\mathcal{O} \subset M$. We know that $\partial\Pi_j \subset \Pi$ has rank 3. Hence $\left[\xi_+, \eta_-\right]$ is a section in Π such that $P_\pm\left[\xi_+, \eta_-\right] \notin C^\infty\left(R^1_\pm\right)$. This means that $C^\infty\left(\Pi^2_+\right)$ and $C^\infty\left(\Pi^2_-\right)$ is locally generated by $\xi_+, P_+\left[\xi_+, \eta_-\right]$ and $\eta_-, P_-\left[\xi_+, \eta_-\right]$ correspondingly. Assume that $\gamma_\pm \in \mathbb{R}\cdot P_\pm\left[\xi_+, \eta_-\right]$, then we can uniquely determine ξ_+, η_-, γ_+ and γ_- by imposing the conditions

$$N_j\left(\xi_+, \gamma_+\right) = \eta_-, \quad N_j\left(\eta_-, \gamma_-\right) = \xi_+, \quad \Omega\left(\xi_+, \gamma_+\right) = 1, \quad \Omega\left(\eta_-, \gamma_-\right) = 1.$$

This system is solvable since Π^2_+ and Π^2_- are orthogonal with respect to N_j. So we have a basis frame on $\mathcal{O}$, namely

$$\left\{\xi_+, \gamma_+, \eta_-, \gamma_-, X_1\right\}.$$

Lemma 4.16 $\sigma\left(\gamma_+\right) = -\frac{1}{2}$, $\sigma\left(\gamma_-\right) = \frac{1}{2}$.

Proof. From theorem 4.8 and the properties of ξ_+, η_-, γ_+ and γ_- we have

$$\begin{aligned} R^\sigma_j\left(\xi_+, \gamma_+\right) &= N_j\left(\xi_+, \gamma_+\right) + \gamma_+\sigma\left(\xi_+\right) - \xi_+\sigma\left(\gamma_+\right) + \gamma_+\sigma\left(\xi_+\right) - \xi_+\sigma\left(\gamma_+\right) \\ &= \eta_- - 2\sigma\left(\gamma_+\right)\xi_+. \end{aligned}$$

We also know

$$R^\sigma_j\left(\xi_+, \gamma_+\right) = -2jX_\sigma \otimes \Omega\left(\xi_+, \gamma_+\right) = -2jX_\sigma.$$

Hence $\eta_- - 2\sigma\left(\gamma_+\right)\xi_+ = -2jX_\sigma$. Similarly for η_-, γ_- we have the two equations

$$\begin{aligned} R^\sigma_j\left(\eta_-, \gamma_-\right) &= N_j\left(\eta_-, \gamma_-\right) - \gamma_-\sigma\left(\eta_-\right) + \eta_-\sigma\left(\gamma_-\right) - \gamma_-\sigma\left(\eta_-\right) + \eta_-\sigma\left(\gamma_-\right) \\ &= \xi_+ + 2\sigma\left(\gamma_-\right)\eta_- \end{aligned}$$

and

$$R^\sigma_j\left(\eta_-, \gamma_-\right) = -2jX_\sigma \otimes \Omega\left(\eta_-, \gamma_-\right) = -2jX_\sigma.$$

Thus $\xi_+ + 2\sigma\left(\gamma_-\right)\eta_- = -2jX_\sigma$. So

$$X_\sigma = \sigma\left(\gamma_+\right)\xi_+ + \frac{1}{2}\eta_- = -\frac{1}{2}\xi_+ + \sigma\left(\gamma_-\right)\eta_-.$$

Since ξ_+ and η_- are linearly independent, we conclude that $\sigma\left(\gamma_+\right)\xi_+ = -\frac{1}{2}\xi_+$ and $\frac{1}{2}\eta_- = \sigma\left(\gamma_-\right)\eta_-$. ■

Definition 4.17 *Let us call a generalized Monge-Ampere equation (ω, θ)* ***nondegenerate*** *in a neighborhood $\mathcal{O}$ in M if:*

- $\operatorname{Im} N_j = \Pi_j$ *is a rank 2 distribution with no singularities within $\mathcal{O}$.*
- Π_j *is nonintegrable and the rank 3 distribution $\partial\Pi_j$ has no singularities within $\mathcal{O}$.*
- *Given generators $\xi_+, \eta_- \in C^\infty\left(R^1_+\right) \times C^\infty\left(R^1_-\right)$ of Π_j, the vector fields ξ_+, η_-, $P_+\left[\xi_+, \eta_-\right]$ and $P_-\left[\xi_+, \eta_-\right]$ are linearly independent in $\mathcal{O}$.*

Theorem 4.18 (Main theorem) *A nondegenerate generalized hyperbolic Monge-Ampere equation (ω, θ) defines a canonical $\{e\}$-structure, i.e. a field of basis frames $\{P_1, P_2, Q_1, Q_2, U\}$. This structure is a complete invariant, i.e. two nondegenerate hyperbolic Monge-Ampere equations are isomorphic if and only if the corresponding $\{e\}$-structures are. The classifying $\{e\}$-structure satisfies the following relations:*

$d\omega\left(\downarrow, \longrightarrow\right)$	P_1	P_2	Q_1	Q_2	U
P_1	0	0	1	0	0
P_2	0	0	0	1	0
Q_1	-1	0	0	0	0
Q_2	0	-1	0	0	0
U	0	0	0	0	0

X	P_1	P_2	Q_1	Q_2	U
$\omega(X)$	0	0	0	0	1
jX	P_1	$-P_2$	Q_1	$-Q_2$	$-$
$\sigma(X)$	0	0	$-\frac{1}{2}$	$\frac{1}{2}$	$-$

$\theta\left(\downarrow, \longrightarrow\right)$	P_1	P_2	Q_1	Q_2
P_1	0	0	1	0
P_2	0	0	0	-1
Q_1	-1	0	0	0
Q_2	0	1	0	0

$N_j\left(\downarrow, \longrightarrow\right)$	P_1	P_2	Q_1	Q_2
P_1	0	0	P_2	0
P_2	0	0	0	P_1
Q_1	$-P_2$	0	0	0
Q_2	0	$-P_1$	0	0

Proof. The tables for $\omega, d\omega, j, N_j$ and σ follow from the construction of the $\{e\}$-structure described in the beginning of this section. We have just renamed the generating sections:

$$P_1 = \xi_+, \qquad P_2 = \eta_-, \qquad Q_1 = \gamma_+, \qquad Q_2 = \gamma_-, \qquad U = X_1.$$

The relation $\theta(X, Y) = d\omega|_\Pi(jX, Y)$ implies the table for θ.

Given an $\{e\}$-structure, satisfying the tables in the theorem, we define the generalized Monge-Ampere equation (ω, θ). Note that the Nijenhuis tensor given by the table must coincide with the Nijenhuis tensor defined from the almost product structure j given by the table. Equivalently we can require that $d_\Pi\theta$ coincides with $d\omega|_\Pi \wedge \sigma$. Under the

fulfillment of either of these conditions, a given $\{e\}$-structure $\{P_1, P_2, Q_1, Q_2, U\}$ uniquely determines the Monge-Ampere equation. ■

An $\{e\}$-structure, $\{e_i\}_{i=1}^5$, give us (potentially) a huge set of invariants of the equation, namely the set of structure coefficients c_{jk}^i from the decomposition $[e_j, e_k] = \sum_i c_{jk}^i e_i$.

Example 4.19 (Nonlinear 4) *We continue the study of the generalized Monge-Ampere equation*

$$(\omega, \theta) = (du - p_1 dq_1 - p_2 dq_2, dq_1 \wedge dp_1 - dq_2 \wedge dp_2 + S dq_2 \wedge dp_1 - T dp_1 \wedge dp_2)$$

represented on the manifold $M = \mathbb{R}^5(q_1, q_2, u, p_1, p_2)$. *We see from example 3.31 that*

$$\begin{aligned} R_+^1 &= \langle \mathcal{D}_{q_1} \rangle = \left\langle \xi_+ \right\rangle, \\ R_-^1 &= \langle (T\mathcal{D}_{q_1}(S) - S\mathcal{D}_{q_1}(T))\,\mathcal{D}_{q_1} + 2\mathcal{D}_{q_1}(T)\,\mathcal{D}_{q_2} - 2\mathcal{D}_{q_1}(S)\,\partial_{p_2} \rangle = \left\langle \eta_- \right\rangle. \end{aligned}$$

So the commutator $\left[\xi_+, \eta_-\right]$ *becomes*

$$\left[\xi_+, \eta_-\right] = \left(T\mathcal{D}_{q_1}^2(S) - S\mathcal{D}_{q_1}^2(T)\right)\mathcal{D}_{q_1} + 2\mathcal{D}_{q_1}^2(T)\,\mathcal{D}_{q_2} - 2\mathcal{D}_{q_1}^2(S)\,\partial_{p_2},$$

where $\mathcal{D}_{q_1}^2 = \mathcal{D}_{q_1} \circ \mathcal{D}_{q_1}$. *We see that* $\Pi_j = R_+^1 \oplus R_-^1$ *is nonintegrable. From the operator* j *in the previous example we can calculate that*

$$\begin{aligned} P_+\left[\xi_+, \eta_-\right] &= 0, \\ P_-\left[\xi_+, \eta_-\right] &= \left[\xi_+, \eta_-\right]. \end{aligned}$$

Thus (ω, θ) *is degenerate according to our definition.*

Example 4.20 *Consider the Monge-Ampere equation*

$$Q - T\frac{\partial^2 u}{\partial q_1^2} + Q\frac{\partial^2 u}{\partial q_2^2} + 2\frac{\partial^2 u}{\partial q_1 \partial q_2} - T\left(\frac{\partial^2 u}{\partial q_1^2}\frac{\partial^2 u}{\partial q_2^2} - \left(\frac{\partial^2 u}{\partial q_1 \partial q_2}\right)^2\right) = 0.$$

Represented in $\mathbb{R}^5(q_1, q_2, u, p_1, p_2)$ *with the contact form* $\omega = du - p_1 dq_1 - p_2 dq_2$, *this equation corresponds to the effective form*

$$\theta = Q dq_1 \wedge dq_2 + T dq_2 \wedge dp_1 + Q dq_1 \wedge dp_2 + dq_1 \wedge dp_1 - dq_2 \wedge dp_2 - T dp_1 \wedge dp_2.$$

One can check that

$$j = \begin{bmatrix} 1 & T & 0 & T \\ Q & -1 & -T & 0 \\ 0 & Q & 1 & Q \\ -Q & 0 & T & -1 \end{bmatrix}.$$

This leads to the splitting of $\Pi = \langle \mathcal{D}_{q_1}, \mathcal{D}_{q_2}, \partial_{p_1}, \partial_{p_2} \rangle$ *into the sum of*

$$\Pi_+^2 = \langle 2\mathcal{D}_{q_1} + Q\mathcal{D}_{q_2} - Q\partial_{p_2}, T\mathcal{D}_{q_1} + Q\partial_{p_1} \rangle$$

and

$$\Pi_-^2 = \langle \mathcal{D}_{q_2} - \partial_{p_2}, T\mathcal{D}_{q_1} + Q\partial_{p_1} - 2\partial_{p_2} \rangle .$$

Denote

$$\begin{array}{ll} \xi_+ = 2\mathcal{D}_{q_1} + Q\mathcal{D}_{q_2} - Q\partial_{p_2}, & \eta_+ = T\mathcal{D}_{q_1} + Q\partial_{p_1}, \\ \xi_- = \mathcal{D}_{q_2} - \partial_{p_2}, & \eta_- = T\mathcal{D}_{q_1} + Q\partial_{p_1} - 2\partial_{p_2}. \end{array}$$

Then

$$\begin{aligned} N_j\left(\xi_+, \eta_+\right) &= 4P_- \circ P_\chi\left[\xi_+, \eta_+\right] \\ &= \left\{-4Q\mathcal{D}_{q_1}(T) - 2Q^2\mathcal{D}_{q_2}(T) + 2Q^2\partial_{p_2}(T) - 4Q\partial_{p_1}(Q)\right. \\ &\quad \left. +2QT\mathcal{D}_{q_2}(Q) - 2QT\partial_{p_2}(Q)\right\}\left(\mathcal{D}_{q_2} - \partial_{p_2}\right). \end{aligned}$$

Similarly we find that

$$\begin{aligned} N_j\left(\xi_-, \eta_-\right) &= 4P_+ \circ P_\chi\left[\xi_-, \eta_-\right] \\ &= 4\left(\mathcal{D}_{q_2}(T) - \partial_{p_2}(T)\right)\mathcal{D}_{q_1} \\ &\quad +2\left(Q\left(\mathcal{D}_{q_2}(T) - \partial_{p_2}(T)\right) - T\left(\mathcal{D}_{q_2}(Q) - \partial_{p_2}(Q)\right)\right)\mathcal{D}_{q_2} \\ &\quad +4\left(\mathcal{D}_{q_2}(Q) - \partial_{p_2}(Q)\right)\partial_{p_1} \\ &\quad -2\left(Q\left(\mathcal{D}_{q_2}(T) - \partial_{p_2}(T)\right) - T\left(\mathcal{D}_{q_2}(Q) - \partial_{p_2}(Q)\right)\right)\partial_{p_2}. \end{aligned}$$

Let us denote

$$A = \mathcal{D}_{q_2}(T) - \partial_{p_2}(T) \qquad \text{and} \qquad B = \mathcal{D}_{q_2}(Q) - \partial_{p_2}(Q).$$

Then we can write

$$R_+^1 = \langle 2A\mathcal{D}_{q_1} + (QA - TB)\mathcal{D}_{q_2} + 2B\partial_{p_1} - (QA - TB)\partial_{p_2} \rangle = \langle X_+ \rangle$$

and

$$R_-^1 = \langle \mathcal{D}_{q_2} - \partial_{p_2} \rangle = \langle Y_- \rangle .$$

One can check that $\Pi_j = R_+^1 \oplus R_-^1$ *is nonintegrable and that*

$$\begin{aligned} P_+\left[X_+, Y_-\right] &= \left[X_+, Y_-\right], \\ P_-\left[X_+, Y_-\right] &= 0. \end{aligned}$$

So (ω, θ) *is degenerate according to our definition.*

Chapter 5

Appendix A

This appendix contains the proof of lemma 3.20. We should show that $N_j(\xi,\eta)(m) = N_j(\xi(m),\eta(m))$ for all sections ξ,η in Π. Let $e_1,\ldots,e_4$ be generators of $C^\infty(\Pi)$ in a neighborhood $\mathcal{O}_m \subset M$ containing the point m. Assume that $\langle e_1,e_2\rangle = \Pi_+^2$ and $\langle e_3,e_4\rangle = \Pi_-^2$. Consider sections $\xi,\widehat{\xi},\Delta\xi,\eta,\widehat{\eta},\Delta\eta \in C^\infty\left(\Pi_+^2\right)$ such that

$$\widehat{\xi} = \xi + \Delta\xi, \qquad \widehat{\eta} = \eta + \Delta\eta,$$

and

$$\widehat{\xi}(m) = \xi(m), \qquad \widehat{\eta}(m) = \eta(m),$$

i.e. $\Delta\xi(m) = \Delta\eta(m) = 0$. Assume that they decompose like

$$\begin{aligned} \xi &= A_1e_1 + \cdots + A_4e_4, & \Delta\xi &= a_1e_1 + \cdots + a_4e_4, \\ \eta &= B_1e_1 + \cdots + B_4e_4, & \Delta\eta &= b_1e_1 + \cdots + b_4e_4, \end{aligned}$$

where $A_i,a_i,B_i,b_i \in C^\infty(M)$ and $a_i(m) = b_i(m) = 0$ for $i = 1,\ldots,4$. Consider the following difference:

$$\begin{aligned} N_j\left(\widehat{\xi},\widehat{\eta}\right) - N_j(\xi,\eta) &= P_\chi\left(\left[j\widehat{\xi},j\widehat{\eta}\right] - \widetilde{j}\left[j\widehat{\xi},\widehat{\eta}\right] - \widetilde{j}\left[\widehat{\xi},j\widehat{\eta}\right] + \left[\widehat{\xi},\widehat{\eta}\right]\right) \\ &\quad -P_\chi\left([j\xi,j\eta] - \widetilde{j}[j\xi,\eta] - \widetilde{j}[\xi,j\eta] + [\xi,\eta]\right) \\ &= P_\chi\Big([j\xi,j\Delta\eta] - \widetilde{j}[j\xi,\Delta\eta] - \widetilde{j}[\xi,j\Delta\eta] + [\xi,\Delta\eta] \\ &\quad + [j\Delta\xi,j\eta] - \widetilde{j}[j\Delta\xi,\eta] - \widetilde{j}[\Delta\xi,j\eta] + [\Delta\xi,\eta] \\ &\quad + [j\Delta\xi,j\Delta\eta] - \widetilde{j}[j\Delta\xi,\Delta\eta] - \widetilde{j}[\Delta\xi,j\Delta\eta] + [\Delta\xi,\Delta\eta]\Big). \end{aligned}$$

We decompose into generating sections and use that $je_1 = e_1, je_2 = e_2$ while $je_3 = -e_3, je_4 = -e_4$.

$$\begin{aligned}
N_j\left(\widehat{\xi},\widehat{\eta}\right) - N_j\left(\xi,\eta\right) &= \sum_{i=1}^{2}\sum_{k=1}^{2} P_\chi\Big(\left[A_ie_i,b_ke_k\right] - \widetilde{j}\left[A_ie_i,b_ke_k\right] - \widetilde{j}\left[A_ie_i,b_ke_k\right] \\
&\quad + \left[A_ie_i,b_ke_k\right] + \left[a_ie_i,B_ke_k\right] - \widetilde{j}\left[a_ie_i,B_ke_k\right] - \widetilde{j}\left[a_ie_i,B_ke_k\right] \\
&\quad + \left[a_ie_i,B_ke_k\right] + \left[a_ie_i,b_ke_k\right] - \widetilde{j}\left[a_ie_i,b_ke_k\right] - \widetilde{j}\left[a_ie_i,b_ke_k\right] + \left[a_ie_i,b_ke_k\right]\Big) \\
&\quad + \sum_{i=1}^{2}\sum_{k=3}^{4} P_\chi\Big(-\left[A_ie_i,b_ke_k\right] - \widetilde{j}\left[A_ie_i,b_ke_k\right] + \widetilde{j}\left[A_ie_i,b_ke_k\right] \\
&\quad + \left[A_ie_i,b_ke_k\right] - \left[a_ie_i,B_ke_k\right] - \widetilde{j}\left[a_ie_i,B_ke_k\right] + \widetilde{j}\left[a_ie_i,B_ke_k\right] \\
&\quad + \left[a_ie_i,B_ke_k\right] - \left[a_ie_i,b_ke_k\right] - \widetilde{j}\left[a_ie_i,b_ke_k\right] + \widetilde{j}\left[a_ie_i,b_ke_k\right] + \left[a_ie_i,b_ke_k\right]\Big) \\
&\quad + \sum_{i=3}^{4}\sum_{k=1}^{2} P_\chi\Big(-\left[A_ie_i,b_ke_k\right] - \widetilde{j}\left[A_ie_i,b_ke_k\right] + \widetilde{j}\left[A_ie_i,b_ke_k\right] \\
&\quad + \left[A_ie_i,b_ke_k\right] - \left[a_ie_i,B_ke_k\right] - \widetilde{j}\left[a_ie_i,B_ke_k\right] + \widetilde{j}\left[a_ie_i,B_ke_k\right] \\
&\quad + \left[a_ie_i,B_ke_k\right] - \left[a_ie_i,b_ke_k\right] - \widetilde{j}\left[a_ie_i,b_ke_k\right] + \widetilde{j}\left[a_ie_i,b_ke_k\right] + \left[a_ie_i,b_ke_k\right]\Big) \\
&\quad + \sum_{i=3}^{4}\sum_{k=3}^{4} P_\chi\Big(\left[A_ie_i,b_ke_k\right] + \widetilde{j}\left[A_ie_i,b_ke_k\right] + \widetilde{j}\left[A_ie_i,b_ke_k\right] \\
&\quad + \left[A_ie_i,b_ke_k\right] + \left[a_ie_i,B_ke_k\right] + \widetilde{j}\left[a_ie_i,B_ke_k\right] + \widetilde{j}\left[a_ie_i,B_ke_k\right] \\
&\quad + \left[a_ie_i,B_ke_k\right] + \left[a_ie_i,b_ke_k\right] + \widetilde{j}\left[a_ie_i,b_ke_k\right] + \widetilde{j}\left[a_ie_i,b_ke_k\right] + \left[a_ie_i,b_ke_k\right]\Big) \\
&= 2\sum_{i=1}^{2}\sum_{k=1}^{2} P_\chi\Big(\left[A_ie_i,b_ke_k\right] - \widetilde{j}\left[A_ie_i,b_ke_k\right] + \left[a_ie_i,B_ke_k\right] \\
&\quad - \widetilde{j}\left[a_ie_i,B_ke_k\right] + \left[a_ie_i,b_ke_k\right] - \widetilde{j}\left[a_ie_i,b_ke_k\right]\Big) \\
&\quad + 2\sum_{i=3}^{4}\sum_{k=3}^{4} P_\chi\Big(\left[A_ie_i,b_ke_k\right] + \widetilde{j}\left[A_ie_i,b_ke_k\right] + \left[a_ie_i,B_ke_k\right] \\
&\quad + \widetilde{j}\left[a_ie_i,B_ke_k\right] + \left[a_ie_i,b_ke_k\right] + \widetilde{j}\left[a_ie_i,b_ke_k\right]\Big).
\end{aligned}$$

Recall that

$$\left[F\xi,G\eta\right] = FG\left[\xi,\eta\right] + FL_\xi\left(G\right)\eta - GL_\eta\left(F\right)\xi$$

holds for any $F, G \in C^{\infty}(M)$ and $\xi, \eta \in C^{\infty}\Pi$. Then we get

$$\begin{aligned}
N_j\left(\widehat{\xi},\widehat{\eta}\right) - N_j(\xi,\eta) &= 2\sum_{i=1}^{2}\sum_{k=1}^{2} P_{\chi}\left(A_i b_k[e_i,e_k] + A_i L_{e_i}(b_k)\, e_k - b_k L_{e_k}(A_i)\, e_i\right. \\
&\quad -A_i b_k \widetilde{j}[e_i,e_k] - A_i L_{e_i}(b_k)\,\widetilde{j}e_k + b_k L_{e_k}(A_i)\,\widetilde{j}e_i \\
&\quad +a_i B_k[e_i,e_k] + a_i L_{e_i}(B_k)\, e_k - B_k L_{e_k}(a_i)\, e_i \\
&\quad -a_i B_k \widetilde{j}[e_i,e_k] - a_i L_{e_i}(B_k)\,\widetilde{j}e_k + B_k L_{e_k}(a_i)\,\widetilde{j}e_i \\
&\quad +a_i b_k[e_i,e_k] + a_i L_{e_i}(b_k)\, e_k - b_k L_{e_k}(a_i)\, e_i \\
&\quad \left. -a_i b_k \widetilde{j}[e_i,e_k] - a_i L_{e_i}(b_k)\,\widetilde{j}e_k + b_k L_{e_k}(a_i)\,\widetilde{j}e_i\right) \\
&\quad +2\sum_{i=3}^{4}\sum_{k=3}^{4} P_{\chi}\left(A_i b_k[e_i,e_k] + A_i L_{e_i}(b_k)\, e_k - b_k L_{e_k}(A_i)\, e_i\right. \\
&\quad +\widetilde{j}A_i b_k[e_i,e_k] + \widetilde{j}A_i L_{e_i}(b_k)\, e_k - \widetilde{j}b_k L_{e_k}(A_i)\, e_i \\
&\quad +a_i B_k[e_i,e_k] + a_i L_{e_i}(B_k)\, e_k - B_k L_{e_k}(a_i)\, e_i \\
&\quad +\widetilde{j}a_i B_k[e_i,e_k] + \widetilde{j}a_i L_{e_i}(B_k)\, e_k - \widetilde{j}B_k L_{e_k}(a_i)\, e_i \\
&\quad +\widetilde{j}a_i b_k[e_i,e_k] + \widetilde{j}a_i L_{e_i}(b_k)\, e_k - \widetilde{j}b_k L_{e_k}(a_i)\, e_i \\
&\quad \left. +a_i b_k[e_i,e_k] + a_i L_{e_i}(b_k)\, e_k - b_k L_{e_k}(a_i)\, e_i\right) \\
&= 2\sum_{i=1}^{2}\sum_{k=1}^{2} P_{\chi}\left(A_i b_k[e_i,e_k] - A_i b_k \widetilde{j}[e_i,e_k] + a_i B_k[e_i,e_k]\right. \\
&\quad \left. -a_i B_k \widetilde{j}[e_i,e_k] + a_i b_k[e_i,e_k] - a_i b_k \widetilde{j}[e_i,e_k]\right) \\
&\quad +2\sum_{i=3}^{4}\sum_{k=3}^{4} P_{\chi}\left(A_i b_k[e_i,e_k] + A_i b_k \widetilde{j}[e_i,e_k] + a_i B_k[e_i,e_k]\right. \\
&\quad \left. +a_i B_k \widetilde{j}[e_i,e_k] + a_i b_k \widetilde{j}[e_i,e_k] + a_i b_k[e_i,e_k]\right).
\end{aligned}$$

Evaluating at the point m we see that

$$N_j\left(\widehat{\xi},\widehat{\eta}\right)(m) - N_j(\xi,\eta)(m) = 0.$$

So we conclude that N_j is a tensor.

Chapter 6

Appendix B

Here we prove proposition 4.4. Let $\xi, \eta, \gamma \in \mathcal{D}_1(M)$. Then we have:

$$\begin{aligned} d\widetilde{\theta}(\xi,\eta,\gamma) &= \left(i_\xi d\widetilde{\theta}\right)(\eta,\gamma) \\ &= \left(L_\xi\widetilde{\theta}\right)(\eta,\gamma) - \left(di_\xi\widetilde{\theta}\right)(\eta,\gamma). \end{aligned}$$

Using

$$L_\xi\left(\widetilde{\theta}(\eta,\gamma)\right) = \left(L_\xi\widetilde{\theta}\right)(\eta,\gamma) + \widetilde{\theta}(L_\xi\eta,\gamma) + \widetilde{\theta}(\eta, L_\xi\gamma),$$

we get that

$$d\widetilde{\theta}(\xi,\eta,\gamma) = L_\xi\left(\widetilde{\theta}(\eta,\gamma)\right) - \widetilde{\theta}(L_\xi\eta,\gamma) - \widetilde{\theta}(\eta, L_\xi\gamma) - \left(di_\xi\widetilde{\theta}\right)(\eta,\gamma).$$

Further

$$\widetilde{\theta} = i_{\widetilde{j}}\widetilde{\Omega} \Longleftrightarrow i_\xi\widetilde{\theta} = i_{\widetilde{j}\xi}\widetilde{\Omega}$$

implies

$$d\widetilde{\theta}(\xi,\eta,\gamma) = L_\xi\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right) - \widetilde{\Omega}\left(\widetilde{j}L_\xi\eta,\gamma\right) - \widetilde{\Omega}\left(\widetilde{j}\eta, L_\xi\gamma\right) - \left(di_{\widetilde{j}\xi}\widetilde{\Omega}\right)(\eta,\gamma).$$

As for $L_\xi\left(\widetilde{\theta}(\eta,\gamma)\right)$, we can write:

$$\begin{aligned} d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right) &= L_{\widetilde{j}\xi}\left(\widetilde{\Omega}(\eta,\gamma)\right) - \widetilde{\Omega}\left(L_{\widetilde{j}\xi}\eta,\gamma\right) - \widetilde{\Omega}\left(\eta, L_{\widetilde{j}\xi}\gamma\right) - \left(di_{\widetilde{j}\xi}\widetilde{\Omega}\right)(\eta,\gamma), \\ d\widetilde{\Omega}\left(\xi,\widetilde{j}\eta,\gamma\right) &= L_\xi\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right) - \widetilde{\Omega}\left(L_\xi\widetilde{j}\eta,\gamma\right) - \widetilde{\Omega}\left(\widetilde{j}\eta, L_\xi\gamma\right) - \left(di_\xi\widetilde{\Omega}\right)\left(\widetilde{j}\eta,\gamma\right). \end{aligned}$$

Using these equations, we get:

$$\begin{aligned} d\widetilde{\theta}\left(\xi,\eta,\gamma\right) &= d\widetilde{\Omega}\left(\xi,\widetilde{j}\eta,\gamma\right)+\widetilde{\Omega}\left(L_{\xi}\widetilde{j}\eta,\gamma\right)+\left(di_{\xi}\widetilde{\Omega}\right)\left(\widetilde{j}\eta,\gamma\right)-\widetilde{\Omega}\left(\widetilde{j}L_{\xi}\eta,\gamma\right) \\ &\quad +d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right)-L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(L_{\widetilde{j}\xi}\eta,\gamma\right)+\widetilde{\Omega}\left(\eta,L_{\widetilde{j}\xi}\gamma\right) \\ &= d\widetilde{\Omega}\left(\xi,\widetilde{j}\eta,\gamma\right)+d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right)+\widetilde{\Omega}\left(\left[\xi,\widetilde{j}\eta\right],\gamma\right)+\left(di_{\xi}\widetilde{\Omega}\right)\left(\widetilde{j}\eta,\gamma\right) \\ &\quad -\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)-L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right)+\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right). \end{aligned}$$

The infinitesimal Stokes formula:

$$di_{\xi}\widetilde{\Omega}=L_{\xi}\widetilde{\Omega}-i_{\xi}d\widetilde{\Omega},$$

gives us:

$$\begin{aligned} d\widetilde{\theta}\left(\xi,\eta,\gamma\right) &= d\widetilde{\Omega}\left(\xi,\widetilde{j}\eta,\gamma\right)+d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right)+\widetilde{\Omega}\left(\left[\xi,\widetilde{j}\eta\right],\gamma\right)+\left(L_{\xi}\widetilde{\Omega}-i_{\xi}d\widetilde{\Omega}\right)\left(\widetilde{j}\eta,\gamma\right) \\ &\quad -\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)-L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right)+\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right) \\ &= d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right)+\left(L_{\xi}\widetilde{\Omega}\right)\left(\widetilde{j}\eta,\gamma\right) \\ &\quad +\widetilde{\Omega}\left(\left[\xi,\widetilde{j}\eta\right],\gamma\right)-\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)-L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right) \\ &\quad +\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right). \end{aligned}$$

We then use that

$$L_{\xi}\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right)=\left(L_{\xi}\widetilde{\Omega}\right)\left(\widetilde{j}\eta,\gamma\right)+\widetilde{\Omega}\left(L_{\xi}\widetilde{j}\eta,\gamma\right)+\widetilde{\Omega}\left(\widetilde{j}\eta,L_{\xi}\gamma\right),$$

and we get:

$$\begin{aligned}
d\widetilde{\theta}\left(\xi,\eta,\gamma\right) &= d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right)+L_{\xi}\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right)-\widetilde{\Omega}\left(L_{\xi}\widetilde{j}\eta,\gamma\right)\\
&\quad -\widetilde{\Omega}\left(\widetilde{j}\eta,L_{\xi}\gamma\right)+\widetilde{\Omega}\left(\left[\xi,\widetilde{j}\eta\right],\gamma\right)-\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)\\
&\quad -L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right)+\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right)\\
&= +d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right)+L_{\xi}\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right)-\widetilde{\Omega}\left(\left[\xi,\widetilde{j}\eta\right],\gamma\right)\\
&\quad -\widetilde{\Omega}\left(\widetilde{j}\eta,\left[\xi,\gamma\right]\right)+\widetilde{\Omega}\left(\left[\xi,\widetilde{j}\eta\right],\gamma\right)-\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)\\
&\quad -L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right)+\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right)\\
&= L_{\xi}\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right)-L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right)-\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)\\
&\quad +\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right)-\widetilde{\Omega}\left(\widetilde{j}\eta,\left[\xi,\gamma\right]\right)+d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right).
\end{aligned}$$

Since $\widetilde{\theta}$ and $\widetilde{\Omega}$ are skew symmetric, we have that:

$$\begin{aligned}
\widetilde{\Omega}\left(\widetilde{j}\eta,\left[\xi,\gamma\right]\right) &= i_{\widetilde{j}}\widetilde{\theta}\left(\eta,\left[\xi,\gamma\right]\right)=-i_{\widetilde{j}}\widetilde{\theta}\left(\left[\xi,\gamma\right],\eta\right)=-\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\gamma\right],\eta\right)\\
&= \widetilde{\Omega}\left(\eta,\widetilde{j}\left[\xi,\gamma\right]\right).
\end{aligned}$$

So finally we get:

$$\begin{aligned}
d\widetilde{\theta}\left(\xi,\eta,\gamma\right) &= L_{\xi}\left(\widetilde{\Omega}\left(\widetilde{j}\eta,\gamma\right)\right)-L_{\widetilde{j}\xi}\left(\widetilde{\Omega}\left(\eta,\gamma\right)\right)+\widetilde{\Omega}\left(\left[\widetilde{j}\xi,\eta\right],\gamma\right)\\
&\quad -\widetilde{\Omega}\left(\widetilde{j}\left[\xi,\eta\right],\gamma\right)+\widetilde{\Omega}\left(\eta,\left[\widetilde{j}\xi,\gamma\right]\right)-\widetilde{\Omega}\left(\eta,\widetilde{j}\left[\xi,\gamma\right]\right)\\
&\quad +d\widetilde{\Omega}\left(\widetilde{j}\xi,\eta,\gamma\right),
\end{aligned}$$

just as we should prove.

Bibliography

[G] E. Goursat, *Leçons sur l'intégration des équations aux dérivées partielles du second ordre à deux variables independantes*, tome I, Hermann, Paris, 1896.

[KLV] I. Krasil'shchik, V. Lychagin, A. Vinigradov, *Geometry of jet spaces and nonlinear partial differential equations*, Adv. Stud. Contemp. Math. 1, Gordon and Breach, New York, 1986.

[Kr1] B. S. Kruglikov, *Some classificational problems in four dimensional geometry: distributions, almost complex structures and Monge-Ampère equations*, (in Russian), Mat. Sb. 189 no. 11 (1998), 61-74; e-print: http://xxx.lanl.gov/abs/dg-ga/9611005.

[Kr2] B. Kruglikov, *Classification of Monge-Ampère equations with two variables*, Banach Center Publ. vol. 50 (1999), 179-194.

[Lie] S. Lie, *Gesammelte Abhandlungen*, Band VI, Aschehoug, Oslo, and Teubner, Leipzig, 1922.

[Ly1] V. V. Lychagin, *Contact geometry and nonlinear second order differential equations*, Uspekhi Mat. Nauk 34 no. 1 (1979), 137-165 (in Russian); English transl.: Russian Math. Surveys 34 (1979), 149-180.

[Ly2] V. Lychagin, *Lectures on Geometry of Differential Equations*, Ciclo di conferenze tenute presso il Dipartimento di Metodi e Modelli Matematici per le Scienze Applicate Università "La Sapienza", Roma, 1992.

[LRC] V. V. Lychagin, V, N, Rubtsov, I, V, Chekalov, *A classification of Monge-Ampère equations*, Ann. Sci. Ècole Norm. Sup. (4) 26 (1993), 281-308.

[S] S. Sternberg, *Lectures on Differential Geometry*, Prentice-Hall, Englewood Cliffs, 1964.